



**HAL**  
open science

## 5-Hydroxymethylfurfural (HMF) in Organic Synthesis: A Review of its Recent Applications Towards Fine Chemicals

Weigang Fan, Charlie Verrier, Yves Queneau, Florence Popowycz

► **To cite this version:**

Weigang Fan, Charlie Verrier, Yves Queneau, Florence Popowycz. 5-Hydroxymethylfurfural (HMF) in Organic Synthesis: A Review of its Recent Applications Towards Fine Chemicals. *Current Organic Synthesis*, 2019, 16 (4), pp.583-614. 10.2174/1570179416666190412164738 . hal-02193105

**HAL Id: hal-02193105**

**<https://udl.hal.science/hal-02193105>**

Submitted on 10 Jul 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 5-Hydroxymethylfurfural (HMF) in organic synthesis: a review of its recent applications towards fine chemicals

Weigang Fan, Charlie Verrier,\* Yves Queneau, Florence Popowycz\*  
Université de Lyon, UMR 5246, CNRS, Université Lyon 1, INSA Lyon, CPE Lyon, ICBMS,  
Bâtiment Edgar Lederer, F-69622 Villeurbanne Cedex, France.  
E-mail: charlie.verrier@insa-lyon.fr; florence.popowycz@insa-lyon.fr

**Abstract:** 5-Hydroxymethylfurfural (HMF) is a platform molecule produced from renewable carbohydrate sources such as fructose, glucose, sucrose, cellulose or inulin. Its interest as building block for accessing chemicals of industrial relevance such as monomers or biofuels has resulted in intense research on cleaner and more efficient processes for its manufacture. Its increased availability has also attracted many studies exploring its use towards value-added fine chemicals, thus widening the scope of its applications. The aim of this review is to bring an updated overview on the use of 5-HMF in fine organic synthesis, to the exclusion of already well documented platform molecules and intermediates such as 2,5-diformylfuran, 2,5-furandicarboxylic acid, 2,5-dihydroxymethylfuran, 2,5-dimethylfuran and so on.

## Table of contents

<b>1. Introduction</b>	1
<b>2. Reactions on the carbonyl group</b>	2
<b>2.1 Carbon-carbon bond formation reactions</b>	2
2.1.1 Formation of C-C bonds	3
2.1.2 Formation of C=C bonds	4
2.1.3 Formation of C≡C bonds	10
2.1.4 Oxidative transformations	10
<b>2.2 Carbon-nitrogen bond formation reactions</b>	11
2.2.1 Synthesis of imines, hydrazones, oximes, nitrones and their applications	11
2.2.2 Preparation of amines	13
2.2.3 Synthesis of nitrogen-containing heterocycles	14
2.2.4 Oxidative transformations	16
<b>2.3 Carbon-oxygen and carbon-sulfur bond formation reactions</b>	16
<b>3 Reactions on the hydroxyl group</b>	17
<b>3.1 Conversion to halides</b>	17
<b>3.2 Conversion to ethers and acetals</b>	18
<b>3.3 Esterifications</b>	20
<b>3.4 Amination reactions</b>	21
<b>3.5 Construction of a C-C(Ar) bonds</b>	22
<b>4 Reactions on the furan ring</b>	22
<b>4.1 Ring opening/rearrangement</b>	22
<b>4.2 Singlet oxygen oxidation</b>	24
<b>4.3 Diels-Alder reactions</b>	24
<b>5 Multi-step and multi-functionalization sequences</b>	25
<b>6 Conclusion</b>	28
<b>7 Acknowledgements</b>	28
<b>8 References</b>	28

## 1. Introduction


The exploitation of biomass-derived platform molecules is an increasingly developed strategy for replacing building blocks arising from petrochemicals. However, their use can be limited due to low-tonnage productions and rather high prices. In the past decades, the obvious interest of bio-based platform molecules for developing

novel chemicals, notably in the field of polymer chemistry, has generated intensive efforts for improving their manufacture processes, therefore increasing their availability due to higher scale industrial production and lower cost. A consequence of this increased availability has been a renewed interest of the chemical community to apply these bio-based building blocks in strategies towards fine chemicals also.

Among platform molecules, 5-hydroxymethylfurfural (5-HMF) appears as one of the most promising. This apparently simple molecule, obtained by dehydration of hexoses (as monosaccharides or included in oligo- or polysaccharides), incorporates an aldehyde group at the C2 position, a hydroxymethyl group at the C5 position, as well as a reactive furan ring, offering a wide range of possible functionalizations and chemical transformations. Whereas the first reports related to the formation and uses of 5-HMF appeared at the end of the 19<sup>th</sup> century, it is only in the past decades that this area has entered a boom stage, consistently with the increasing awareness of environmental concerns in the chemical community and the development of biobased chemistry. A specific character of HMF is its relative limited stability, which is a drawback both at the level of its manufacture and of its utilizations. This sensitivity to both heat and acids leads to the occurrence of degradation and undesired reactions such as dimerization or oligomerization.<sup>[1-3]</sup> Taking into account the high potential of 5-HMF, some methods aiming at improving its stability start to appear, such as the recently reported use of sodium dithionite.<sup>[4]</sup>

Several comprehensive recent reviews by Afonso,<sup>[5]</sup> de Vries<sup>[6-7]</sup> and Ananikov,<sup>[8]</sup> extensively cover the synthesis and manufacture of HMF, and/or its catalytic conversion to simple molecules such as 2,5-diformylfuran (DFF) or 2,5-difurandicarboxylic acid (FDCA). They illustrate most immediate industrial relevance and economic interests of HMF and, in a less detailed manner, its applications towards fine chemicals. The purpose of the present review article, as a complementary account to the extensive reviews previously published, is to provide an exhaustive and updated overview describing the synthetic methodologies developed around 5-HMF chemistry, in the context of fine chemicals synthesis, including multi-step and total synthesis. It illustrates the importance of the HMF scaffold as a building block in all areas of synthetic organic chemistry.

The review is organized with respect to the different chemical functions present in HMF and their reactivity. First, transformations of the aldehydic moiety of 5-HMF will be discussed, then the transformations of the hydroxymethyl group, and the modifications of the furan ring (Scheme 1). Finally, multistep and multi-functionalization sequences involving 5-HMF are shown, giving a complete exemplification of the incorporation of this platform molecule into complex chemical architectures.



**Scheme 1.** Overview of 5-HMF chemistry

## 2. Reactions on the carbonyl group

Aldehydes are essential building blocks in organic synthesis. Their high electrophilicity makes them appropriate partners for reactions with a variety of nucleophiles, allowing the formation of single, double or triple carbon-carbon bonds, but also the construction of carbon-heteroatom bonds through condensation reactions. Consequently, most of the direct transformations of 5-HMF toward high value-added chemicals take advantage of its aromatic aldehyde group.


### 2.1 Carbon-carbon bond formation reactions

Carbon-carbon bonds are among the strongest bonds of organic chemistry, and their construction constitutes the basis of most synthetic routes toward organic molecules. Many strategies have been developed to generate new C-C bonds using the aldehydic moiety of 5-HMF, offering accesses to new bio-based compounds with various properties.

### 2.1.1 Formation of C-C bonds


The benzoin condensation is an efficient method to upgrade HMF to C<sub>12</sub> furoins as potential fuel intermediates. During the course of a study on the degradation of HMF in ionic liquids such as 1-ethyl-3-methylimidazolium acetate ([EMIM]OAc), Chen found that 1-ethyl-3-methylimidazolin-2-ylidene carbene could catalyze the self-condensation of HMF to form 5,5'-dihydroxymethylfuroin **1** with high conversion and selectivity (Table 1).<sup>[9]</sup> Since this pioneering work, several NHC catalysts derived from ILs and discrete catalysts proved their efficiency for the benzoin condensation of HMF.<sup>[10-14]</sup> Benzaldehyde lyase could also be used as a biocatalyst for this umpolung condensation.<sup>[15]</sup> However, under those conditions, a partial spontaneous oxidation to the corresponding diketone was observed.

**Table 1.** Organocatalysts and biocatalyst for HMF benzoin condensation




Entry	Catalyst	Conditions	Yield	Ref
1		[EMIM]OAc, 80 °C or 5 mol% [EMIM]Cl, DBU, THF, 80 °C	50% (isolated) or 97% (HPLC)	[9]
2	 TPT	0.1-5 mol% TPT, neat or in THF, 60 °C	87-95% (isolated)	[9-11]
3		10 mol% Silica-g-[BI]-C <sub>12</sub> azolium salt, DBU, THF, 25 °C, 6 h	94% (NMR)	[12]
4		Thiazolium derived ILs, Et <sub>3</sub> N, 120 °C	97-98% (NMR)	[13-14]
5	Benzaldehyde lyase	potassium phosphate buffer /DMSO, rt	70% conversion	[15]

Afonso and coworkers reported the preparation of a highly functionalized furan skeleton *via* a novel reaction pathway from 5-HMF or *O*-protected HMF and a secondary amine.<sup>[16]</sup> HMF could react with morpholine to provide the iminium intermediate **2**, which is in equilibrium with the trienamine **3** under acidic catalysis (Scheme 2). The two intermediates reacted together, releasing a molecule of amine to yield the coupling product **4**.




**Scheme 2.** Homo-bis-vinyllogous Mannich-type reaction on 5-HMF

The synthesis of the symmetric triarylmethane **5** was achieved by reaction of a secondary aniline on 5-HMF in the presence of Yb(OTf)<sub>3</sub> as acidic catalyst, following a double Friedel-Crafts pathway (Scheme 3).<sup>[17]</sup> The scope of the reaction was successfully extended to *O*-substituted hydroxymethylfurfurals. The reaction was found to be significantly accelerated by applying high pressure, especially in the case of less reactive anilines. The authors considered that the first Friedel-Crafts reaction proceeded first *via* the addition of the aniline onto the iminium ion formed by condensation of the secondary aniline on the aldehyde. Then a second Friedel-Crafts reaction took place in a non-concerted mechanism suggesting a secondary carbocation as the transient intermediate for this step.



**Scheme 3.** Synthesis of symmetric triarylmethanes *via* Friedel-Crafts mechanism

A last type of C-C bond forming reaction with carbonyl derivatives exploiting the Morita-Baylis-Hillman reaction has been reported. The use of electron-deficient alkenes such as methyl acrylate<sup>[18]</sup> and acrylamide<sup>[19]</sup> in combination with DABCO in a dioxane/water mixture at room temperature gave access to the conjugated compounds **6** (Scheme 4). Our group more recently extended this reactivity to other solvent mixtures, including bio-based solvent, and to glycosylated derivatives of HMF.<sup>[20-21]</sup>




**Scheme 4.** Morita-Baylis-Hillman reaction of 5-HMF

### 2.1.2 Formation of C=C bonds

Aldol- and Knoevenagel type condensations are classical transformations of the chemistry of the carbonyl group. Aromatic aldehydes, such as HMF, are very good substrates for such reactions, due to the absence of proton on the carbon adjacent to the carbonyl (no competition possibly arising from self-condensation).

Different research groups have explored the reaction of acetone with 5-HMF under basic conditions. This transformation provides a straightforward access to the natural products rhemanones A, B and C, owning interesting biological activities such as blood platelet aggregation inhibitors or immunostimulant. The challenge relies on the control of the mono- or bis-condensation of the ketone on 5-HMF, as the monoadduct leads to rhemanone B and C, whereas bis-condensation would give an advanced intermediate for the preparation of rhemanone A (Table 2). The group of Tamariz used sodium hydroxide to promote the reaction, and a good control of the selectivity (ratio **7** vs **8**) was obtained by tuning the quantity of acetone used.<sup>[22]</sup> The yield of mono adduct **7** reached 91% when acetone was used in excess, whereas the di-adduct **8** was obtained in a 60 % yield when HMF was used in excess. Palkovits described a quite selective mono-condensation mediated by mesoporous magnesium aluminate as the base.<sup>[23]</sup> Roman-Leshkov reported the use of hafnium- $\beta$  zeolite providing compound **7** with >99% selectivity and 73% conversion.<sup>[24]</sup> Another selective mono-condensation, promoted by CO<sub>2</sub> under elevated pressure with a high yield, was reported by Jessop.<sup>[25]</sup> A similar yield of the mono-adduct **7** could be obtained under heterogeneous catalysis conditions using amine-grafted Faujasite (FAU) zeolite nanosheets (Table 2).<sup>[26]</sup> Zirconium carbonate was also used to promote the same condensation under mild conditions.<sup>[27]</sup>


**Table 2.** Synthesis of the precursors of rhemanones



Entry	Conditions	Product (yield)	Ref
1	Acetone (2 eq.), NaOH (aq), EtOH, rt	<b>7</b> (91%)	[22]
2	Acetone (0.5 eq.), NaOH (aq), EtOH, rt	<b>8</b> (60%)	[22]


3	Acetone (as solvent), MgAl <sub>2</sub> O <sub>4</sub> , 140 °C	<b>7</b> (81%) + <b>8</b> (11%)	[23]
4	Acetone (3 eq.), Hf-Beta zeolite, toluene, 90 °C	<b>7</b> (73% conv)	[24]
5	Acetone (solvent), H <sub>2</sub> O, 2.5 MPa CO <sub>2</sub> , 200 °C, 20h	<b>7</b> (95%)	[25]
6	Acetone (solvent), amine-grafted FAU nanosheets, 130 °C	<b>7</b> (92% conv)	[26]
7	Acetone (20 eq.), zirconium carbonate, H <sub>2</sub> O, 54 °C	<b>7</b> (92%)	[27]

Skowronski<sup>[28]</sup> and Corma<sup>[29]</sup> independently reported the efficient condensation of acetophenone derivatives with HMF. The reaction, promoted by potassium hydroxide in methanol, produced 5-hydroxymethyl-furfurylidene-acetophenone **9**, later named furanochalcone, in 82% yield (Scheme 5). The same transformation was also performed with almost quantitative yield in the presence of a catalytic amount of Mg-Al mixed oxide (HTc). Different 5-hydroxymethyl-furanochalcones have then been prepared following similar strategies, and evaluated for diverse biological activities such as phosphatase inhibitor<sup>[30]</sup> or synergistic antifungal.<sup>[31]</sup> The group of Surwayanshi described the preparation of enone **11** by condensation of the dihydro- $\alpha$ -ionone **10** with HMF under microwave irradiation using KF and aluminate as heterogeneous base.<sup>[32]</sup> The product **11** was then evaluated for its antileishmanial properties. A condensation with 1-C-( $\beta$ -D-galactopyranosyl)-propan-2-one in the presence of lithium hydroxide was performed to prepare galactose derivative **12** in 65% yield, which proved to be an effective antimycobacterial agent (Scheme 5).<sup>[33]</sup> A C-4 epimer of **12** could also be prepared from 1-C- $\beta$ -glucosylacetone in dichloromethane in the presence of pyrrolidine in 87% yield.<sup>[34]</sup>




**Scheme 5.** Condensation with ketones

Some more elaborated ketones have been employed in aldol condensations with 5-HMF to produce molecules with high added-value. One example relies on its condensation with cyclooctanone to generate the intermediate **13**, which can be further converted into the thieno[2,3-*b*]pyridine **14**, found to be active against the elongation factor-2 kinase (eEF2-K) (Scheme 6).<sup>[35]</sup> Some benzofuran-3-ones were also condensed with 5-HMF to prepare furanic analogues **15** of auroene. The KOH-mediated aldol condensation of di-hydroxybenzofuranone with HMF provided **15** in 53% yield.<sup>[36]</sup> The microwave activation carried out in a deep eutectic solvent did not bring any improvement in terms of yield.<sup>[37]</sup> Finally, a base-catalyzed aldol condensation between dihydrolevoglucosenone and HMF was described by Witczak and Bielski.<sup>[38]</sup> Piperidine gave superior efficiency when the reaction was conducted in refluxing ethanol, providing the chiral building block **16** in 86% yield. The base-mediated double condensation of cyclopentanone on 5-HMF proved to be an efficient route toward molecule **17**. The latter has been studied as a synergistic antifungal compound,<sup>[39]</sup> but also constituted a valuable intermediate in the preparation of long-chain cycloalkanes from bio-renewable resources.<sup>[40]</sup>




**Scheme 6.** Condensation on more elaborated ketones as targets in medicinal chemistry

The condensation of unsymmetrically substituted ketones bearing two enolizable positions with 5-HMF raises a problem of regioselectivity. Aqueous sodium hydroxide was employed by Amarasekara and coworkers to condense levulinic acid on 5-HMF.<sup>[41]</sup> The reaction produced a mixture of kinetic and thermodynamic products (respectively **18** and **19**) in a 2.5:1 ratio and 82% total yield (Scheme 7). This problem could be avoided by performing the reaction under pyridine/DCM conditions.<sup>[34]</sup> The condensation of 5-HMF with methyl isobutyl ketone (MIBK) in the presence of NaOH, led selectively to the corresponding kinetic enone in 95% GC yield.<sup>[42]</sup>




**Scheme 7.** Regioselective condensation on levulinic acid

The regioselective condensation of acetylacetone with 5-HMF was also studied. Acetylacetone bearing highly acidic protons at C3 position, the selective condensation at C1 and/or C5 without the formation of the Knoevenagel adduct (C3 condensation) represented a synthetic challenge and could lead to curcumin analogs (Scheme 8). A solution emerged from Lee's group<sup>[43]</sup> through a double C1 and C5 enolization in the presence of boric anhydride, tributylborate, butylamine and an excess of 5-HMF, giving the symmetrical double condensation product **20** in 12% yield. When the same reaction was carried out with an excess of diketone, the Knoevenagel adduct **21**, arising from the functionalization at C3, was obtained in 51% yield (Scheme 8).<sup>[44]</sup> The most efficient alternative reported to date for the synthesis of unsymmetrical HMF-derived curcumin analogues such as **22**, consisted in a solid-phase synthesis strategy. Supporting HMF on a 2-chlorotrityl resin led to a significant improvement of the yield.<sup>[45]</sup> After completion of the reaction, the resin was cleaved by treatment with a mixture of DCM/TFA/MeOH, releasing the final product **22** in excellent yield (Scheme 8).




**Scheme 8.** Strategies for regioselective aldolisation of HMF with acetylacetone

The cyanoamide **23** was reacted with HMF to produce the cyano-acrylamide **24** under microwave irradiation in a modest 45% yield (Scheme 9).<sup>[46]</sup> McCluskey reported the reaction of dichlorophenylacetonitrile with 5-HMF under phase transfer catalysis which provided the acrylonitrile **25** in 78% yield.<sup>[47]</sup> In parallel to compound **25**, evaluated for its antitumoral properties, the two other derivatives **26** and **27** (supposed to exhibit significant anti-senile dementia activities) were prepared by condensation with the corresponding activated methylene substrates (Scheme 9).<sup>[48]</sup>



**Scheme 9.** Condensation of cyanoamides and nitriles derivatives with 5-HMF

Cyclic amides such as rhodanine **28**<sup>[49]</sup> and thiohydantoin **30**<sup>[50]</sup> have been employed for the preparation of 5-hydroxymethylfurfurylidene-substituted heterocycles, designed for pharmacological tests (Scheme 10). The furfurylidene-rhodanine **29** was obtained in the presence of ethylenediammonium diacetate with 73% yield whereas the thiohydantoin **31** was formed in the presence of pyrrolidine and indium triflate as Lewis acid.




**Scheme 10.** Cyclic amides in the condensation with 5-HMF

Several activated methylene compounds have been used in Knoevenagel type condensations with HMF yielding push-pull olefins which could be further used as building blocks or directly studied for their biological properties. Highly acidic cyclic 1,3-dicarbonyl derivatives such as Meldrum's acid and barbituric acid proved their efficiency, reacting with HMF smoothly. The condensation with Meldrum's acid using a mixture of piperidine and acetic acid in benzene afforded compound **32** in 72% yield, and its antitumoral and psychotropic activities




were evaluated (Scheme 11).<sup>[51]</sup> Another experimental procedure reported by Afonso referred to the use of water with 1 wt% Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> leading to 96% of yield of the same compound **32**.<sup>[4]</sup> The authors also reported an interesting transformation of **32** to lactone-fused cyclopentenones by treatment with various amines.<sup>[52]</sup> Nikolov carried out the condensation between HMF and barbituric acid in refluxing water, and assessed the photochemical properties of the product **33**.<sup>[53]</sup> Finally, 4-hydroxycoumarin reacted in catalyst-free conditions to produce the dimeric compound **34** in 55% yield.<sup>[54]</sup> The mono-addition led to a Knoevenagel intermediate exhibiting structural analogy with the *ortho*-quinone methide skeleton, facilitating the addition of a second molecule of 4-hydroxycoumarin (Scheme 11).




**Scheme 11.** Condensation of cyclic activated methylene derivatives with 5-HMF

Acyclic activated methylene compounds are also possible nucleophiles for condensation with 5-HMF. For example, the reaction of 5-HMF with malonic acid in the presence of ammonium bicarbonate without solvent at 50 °C afforded the product **35** in 93% yield.<sup>[55]</sup> Decarboxylation took place at a higher temperature in pyridine, leading to the acrylic acid **36** up to 88% yield.<sup>[28, 34]</sup> Finally, Knoevenagel condensation of the dione **37** promoted by piperidine allowed the preparation of the dihydrochromone **38** which could be rapidly converted into the chromone **39**, a photo-activable fluorophore (Scheme 12).<sup>[56]</sup>




**Scheme 12.** Condensation of acyclic activated methylene compounds

Compound **41** was identified as a side product of the reaction between D-glucose and L-lysine under acidic conditions. In order to support the mechanism explaining its formation, a larger amount of compound **41** was successfully prepared by reaction of 5-HMF on the cyclic imine **40** (obtained from piperidine, *N*-chlorosuccinimide and potassium hydroxide) in 64% yield (Scheme 13).<sup>[57]</sup>




**Scheme 13.** Condensation of 5-HMF with compound **40**

Condensation of the diamionitroalkene **42** on 5-HMF provided the new neonicotinoid **43** with insecticidal potency against *Nephotettix bipunctatus*, higher than imidacloprid (Scheme 14).<sup>[58]</sup>




**Scheme 14.** Condensation of 5-HMF with enediamine **42**

The HMF-modified boron-dipyrromethene (BODIPY) **44** was prepared *via* an aldol-like condensation between HMF and 1,3-dimethyl-BODIPY in the presence of pyrrolidine under microwave irradiation (Scheme 15).<sup>[59]</sup>




**Scheme 15.** Synthesis of a BODIPY molecule for bio-imaging application

The indene-derivative **45** was prepared as a mixture of isomers by the condensation of 2-methylindene-3-acetic acid with HMF<sup>[60]</sup>, and its sodium salt displayed anti-proliferative activities (Scheme 16).




**Scheme 16.** Condensation with 2-methylindene-3-acetic acid

In 1984, the first Horner-Wadsworth-Emmons reaction of HMF was reported by Delmas. Reaction with triethylphosphonoacetate using potassium carbonate in a mixture of 1,4-dioxane and water afforded  $\alpha,\beta$ -unsaturated ester **46** as *E* isomer exclusively in 90% yield (Scheme 17).<sup>[61]</sup> When the reaction was performed in alcoholic medium such as *n*-octanol, further transesterification occurred.<sup>[62]</sup> A DBU-mediated Horner-Wadsworth-Emmons reaction on HMF was performed to synthesize  $\alpha,\beta$ -unsaturated imide **47**, which was obtained as a single diastereoisomer in 87% yield, and might further serve as a precursor of  $\beta$ -amino acids.<sup>[63]</sup> The Horner-Wadsworth-Emmons reaction of HMF was also reported as the key step in a one-pot preparation of substituted pyrazoles.<sup>[64]</sup> HMF or *O*-protected derivatives reacted with diethoxyphosphorylacetaldehyde tosylhydrazone in the presence of NaH in THF, yielding  $\alpha,\beta$ -unsaturated tosylhydrazones sodium salts **48**, which in refluxing THF provided the pyrazoles **49** in 60-78% yield for 2 steps (Scheme 17).




**Scheme 17.** Horner-Wadsworth-Emmons reaction on 5-HMF

The  $\alpha,\beta$ -unsaturated ester **46** could also be prepared by one-pot Wittig reaction. The *in situ* formed Wittig reagent reacted with 5-HMF in the presence of tetra-*n*-butylammonium fluoride (TBAF) providing **46** with 81% yield and 85% selectivity in favor of *E* isomer (Scheme 18).<sup>[65]</sup>



**Scheme 18.** One-pot Wittig reaction on 5-HMF


The monomer 5-hydroxymethyl-2-vinylfuran (HMVF) was synthesized using methyltriphenylphosphonium bromide and potassium carbonate in 67% yield, as an adaptation of a procedure used on furfural (Scheme 19).<sup>[66]</sup> The resulting HMVF could be efficiently polymerized by AIBN-initiated radical polymerization. Recently, Ji demonstrated that HMVF itself exhibits strong adhesion to metals *via in-situ* polymerization and crosslinking after either heating or acid treatment, concluding that HMVF could serve as a bio-based adhesive.<sup>[67]</sup>



**Scheme 19.** Direct alkenylation of 5-HMF

### 2.1.3 Formation of C≡C bonds


Direct alkynylation of 5-HMF was performed in 97% yield using Ohira–Bestmann reagent in MeOH in the presence of K<sub>2</sub>CO<sub>3</sub> (Scheme 20). 2-Hydroxymethyl-5-ethynylfuran **50** was used as a platform to conduct various chemical transformations targeting furanic pharmaceuticals as well as conjugated polymers.<sup>[68]</sup>



**Scheme 20.** Direct alkynylation of 5-HMF


### 2.1.4 Oxidative transformations

Van der Eycken and Noël reported an acylation protocol at C2 of *N*-pyrimidylindoles using a range of aldehydes as acylating reagents *via* dual photoredox/transition-metal catalysis in flow conditions (Scheme 21).<sup>[69]</sup> In this work, HMF also showed certain reactivity leading to compound **51**, but with a low isolated yield of 34% in comparison with furfural (85%). This important difference of yield between similar substrates suggests detrimental effect of the hydroxymethyl group, moderately compatible with oxidative SET pathways.




**Scheme 21.** Acylation of indole using HMF in a photoredox strategy

Following a Büchner-Curtius-Slotterbeck reaction, HMF was reacted with diazomethane yielding 1-(5-(hydroxymethyl)furan-2-yl)ethan-1-one (**52**) in 40% yield after 14 days (Scheme 22).<sup>[70]</sup>




**Scheme 22.** Reaction between 5-HMF and diazomethane

A ruthenium dihydride complex was employed as a catalyst for the coupling of 5-HMF and isoprene on both C2 and C5 positions leading to the bis-ketofurane **53** in 48% yield (Scheme 23).<sup>[71]</sup>



**Scheme 23.** Coupling of 5-HMF and isoprene

Das reported a novel route to prepare  $\alpha$ -diketone by performing the NHC-catalyzed HMF self-condensation in DMSO under  $\text{CO}_2$  atmosphere (Scheme 24).<sup>[72]</sup> DFT calculations lightened the role of  $\text{CO}_2$  in the mechanism, by reacting with the hydroxyl of so formed furoin, producing a carboxylate which is a good leaving group. DMSO was prone to therefore act as an oxidation reagent allowing the access to diketone **54**. The non-symmetrical  $\alpha$ -diketone **55** could also be prepared in good yield employing a 1:1.5 ratio of furfural and HMF, although the formation of the homo  $\alpha$ -diketone could not be suppressed completely.




**Scheme 24.** NHC and  $\text{CO}_2$ -mediated homo and cross-coupling of HMF

## 2.2 Carbon-nitrogen bond formation reactions

The condensation of nitrogen nucleophiles on the aldehyde of HMF is a straightforward way to produce a variety of new molecules with interesting properties, but can also be a step of more complex chemical sequences toward original nitrogen-containing scaffolds, useful for agrochemical and pharmaceutical industries.


### 2.2.1 Synthesis of imines, hydrazones, oximes, nitrones and their applications

In the early stage study on HMF from the end of the 19<sup>th</sup> century to the beginning of the 20<sup>th</sup> century, researchers already showed that HMF could react with a range of nitrogen nucleophiles, such as hydroxylamine, arylhydrazine, semioxamazine, semicarbazide and amine to form respectively oxime **56**, hydrazone **57**, semioxamzone **58**, semicarbazone **59** and imine **60**<sup>[73]</sup> (Scheme 25).<sup>[74]</sup> These reactions with nitrogen nucleophiles were carried out mainly for purification and identification purposes because the products are often crystalline and easy to handle.




**Scheme 25.** Reactions between HMF and diverse nitrogen-nucleophiles in early studies

The imines **61** obtained from HMF were subjected to TFA-catalyzed Pudovik reaction to afford the  $\alpha$ -aminophosphonates **62** (Scheme 26).<sup>[75]</sup> In 2018, our group reinvestigated the reaction conditions in a direct 3-CR strategy between 5-HMF, amine and dialkylphosphite. This one-pot approach named Kabachnik-Fields reaction was efficient in the presence of mild Lewis acid catalyst  $\text{I}_2$ .<sup>[76]</sup>




**Scheme 26.** Synthesis of  $\alpha$ -aminophosphonates *via* Pudovik or Kabachnik-Fields reactions

Amarasekara reported an interesting application of the HMF-derived oxime **56**.<sup>[77]</sup> Oxidation of **56** in aqueous bleach solution provided the nitrile oxide **63**, which readily underwent 1,3-dipolar cycloaddition with alkenes or alkynes to give the corresponding dihydro-isoxazoles **64** or isoxazoles **65** depending on the nature of the dipolarophiles (Scheme 27).




**Scheme 27.** Application of HMF-derived oxime in 1,3-dipolar cycloaddition

The condensation of HMF with rhodamine B hydrazone delivered hydrazone **66**, which could serve as a chemosensor for the detection of pH and  $\text{Cu}^{2+}$ .<sup>[78]</sup> A couple of HMF-derived *N*-acyl-hydrazones **67** were prepared as nifuroxazine analogues for antimicrobial activity screening.<sup>[79]</sup> Unfortunately, the HMF-based hydrazones **67** displayed no interesting biological properties. Hydrazone **68** was also prepared to study its activity as a kinase inhibitor,<sup>[80]</sup> whereas compound **69** (from 1-aminoquinolin-2(1*H*)-one) was studied as a ligand for making metal complexes.<sup>[81]</sup>



**Scheme 28.** Examples of valuable HMF-based acylhydrazones and hydrazones


Capon isolated naturally occurring prolinimines from a fish gastrointestinal tract-derived fungus, *Trichoderma sp.* CMB-F563 (Scheme 29).<sup>[82]</sup> To confirm their structures, the authors performed a biomimetic synthesis of the prolinimine **70** by reacting HMF with *N*-amino-L-Pro methyl ester. Further treatment of prolinimine **70** in acidic conditions led to oligomeric prolinimines **71** and **72**.



**Scheme 29.** Synthesis of natural prolinimines from 5-HMF


The preparation of nitrone **73** was investigated by heterogeneous reductive coupling of nitrobenzene and HMF

under H<sub>2</sub> atmosphere in the presence of platinum nanoparticles (Scheme 30).<sup>[83]</sup> The conversion was complete and the selectivity was evaluated at 89% (76% yield) in favor of the nitron (side products expected: partial reduction into imine, total reduction into secondary amine or reduction of 5-HMF into the corresponding symmetrical diol).



**Scheme 30.** Preparation of nitron

The Murai reaction *via* imine directing group assisted C-H activation on the furan ring of HMF derivatives was first reported by Poli, Oble and coworkers (Scheme 31).<sup>[84]</sup> A diamine reacted with the aldehyde group of *O*-protected HMF derivatives to form products **74** in which the amino-imine moiety served as a *N,N'*-bidentate directing group. The reaction was performed using a typical ruthenium catalyst and vinylsilanes as olefin partners in toluene at high temperature. The directing group was removed by hydrolysis during the silica gel column chromatography purification to release the C3-functionalized furfurals **75**. More recently, the same group reported the Ru-catalyzed imine-directed C3-arylation of compound **76** with aryl boronate esters.<sup>[85]</sup> In this case, a *p*-dimethylaminophenylimine proved to be the best directing group. In contrast to their previous study, the arylated imine **77** could be isolated by silica gel chromatography, the arylated aldehydes **78** being further released after hydrolysis with aqueous HCl (Scheme 31).




**Scheme 31.** Imine as directing group for C-H activation

### 2.2.2 Preparation of amines


Many examples of reductive amination of HMF to target 5-aminomethyl-2-furfuryl alcohols have been reported (Table 3). The utilization of NaBH<sub>4</sub> as a reducing agent was proved to be efficient,<sup>[86]</sup> but many studies describe also the use of H<sub>2</sub> together with a wide range of heterogeneous catalysts. Other reducing agents such as water or silanes have also been reported. Throughout these studies, aromatic and aliphatic amines, as well as ammonia, were introduced on the HMF scaffold.

**Table 3.** Reductive amination with simple amines




Entry	Conditions	RNH <sub>2</sub>	Yield	Ref.
1	NaBH <sub>4</sub> , H <sub>2</sub> O or EtOH	aromatic/aliphatic amines	77-99%	[86]
2	H <sub>2</sub> , Raney Ni	NH <sub>3</sub> (liq.)	72%	[87]
3	H <sub>2</sub> (1.2 MPa), Ru(DMP) <sub>2</sub> Cl <sub>2</sub> , EtOH, 60 °C	aromatic/aliphatic amines	66-95%	[88]
4	H <sub>2</sub> (2 MPa), Rh/Al <sub>2</sub> O <sub>3</sub> , 80 °C	NH <sub>3</sub> (aq.)	86%	[89]
5	H <sub>2</sub> (4 MPa), Ru/Nb <sub>2</sub> O <sub>5</sub> , MeOH, 90 °C	NH <sub>3</sub> (in MeOH solution)	96%	[90]
6	H <sub>2</sub> (2 MPa), RuNPs, MeOH, 90 °C	NH <sub>3</sub> (in MeOH solution)	95%	[91]
7	H <sub>2</sub> (4 MPa), Co nanoparticles, <sup>t</sup> BuOH, 120 °C	NH <sub>3</sub> (gas)	89%	[92]
8	H <sub>2</sub> O/CO (2 MPa)/MeOH, Au/TiO <sub>2</sub> -R, 60 °C	aromatic/aliphatic amines	60-99%	[93]

The reductive amination was applied to a phenanthropiperidine derivative using NaBH(OAc)<sub>3</sub> as reducing agent, to deliver the *N*-substituted phenanthropiperidine derivative **79** in 72% yield (Scheme 32). This tylophorine analogue was evaluated with respect to its biological activity and found to inhibit the proliferation of MCF-7 cells but only at micromolar concentration.<sup>[95]</sup> The indolopyrrolocarbazole derivative **80**<sup>[96]</sup> and cytosine derivative **81**<sup>[97]</sup> were also synthesized by reductive amination, and their antitumor effect and DNA methylation inhibition activity were respectively studied (Scheme 32).



**Scheme 32.** Reductive amination with complex amines and derivatives


By switching the ratio of HMF and amine, Zhang et al. were able to convert HMF to bis(hydroxymethylfurfuryl)amines **82** using Ru(DMP)<sub>2</sub>Cl<sub>2</sub> as catalyst (Scheme 33).<sup>[98]</sup> The *in situ* generated secondary amine further reacted with excess of HMF to form an iminium-ion intermediate, followed by hydrogenation to produce bis(hydroxymethylfurfuryl)amines **82**.



**Scheme 33.** Double reductive amination


### 2.2.3 Synthesis of nitrogen-containing heterocycles

The condensation of 5-HMF with 2-aminothiophenol took place smoothly in AcOH giving rise to the benzothiazole **83** quantitatively after spontaneous oxidation (Scheme 34).<sup>[99]</sup> Similarly, the benzo[*d*]imidazole derivative **84**, prepared by reaction of HMF with *o*-phenyldiamine in the presence of DDQ in refluxing EtOH, served as an intermediate for the preparation of molecule **85**, a potential Myc modulator.<sup>[100]</sup>




**Scheme 34.** Synthesis of benzothiazole and benzimidazole derivatives

HMF was often introduced as an aldehyde component in combinatorial chemistry to synthesize libraries of complex heterocyclic derivatives designed for bioactivity screening (Scheme 35). Treatment of HMF with *p*-toluenesulfonylmethyl isocyanide and potassium carbonate in refluxing methanol produced the 5-(5-hydroxymethyl-2-furyl)oxazole **86** in 66% yield.<sup>[101]</sup> In FM-381, an outstanding JAK3 inhibitor, synthesized by Laufer and coworkers,<sup>[102-103]</sup> the furan moiety plays a crucial role with respect to the inhibitory activity, as replacement of the furan link by a phenyl one generally induced lower activities. The 5-HMF was condensed on 4,5-diamino-7-azaindole to provide the tricyclic skeleton **87** as a precursor of FM-381. Sova prepared a range of quinazolinone derivatives such as **88** to study their antimicrobial activities, among which the HMF-derived product has no significant effect.<sup>[104]</sup> A fused tetracyclic heterocycle **89** was prepared in 68% yield by the TFA-catalyzed condensation of HMF and a 2-(1*H*-pyrrol-1-yl)aniline derivative.<sup>[105]</sup> The condensation of HMF with pyrimidinylguanidine furnished the desired pyrimido[1,2-*a*][1,3,5]triazin-6-one **90** in 38% yield.<sup>[106]</sup>



**Scheme 35.** Synthesis of nitrogen-containing heterocycles

An interesting extension of the scope to access more complex molecules was achieved by using multicomponent reactions (Scheme 36). The 3-component condensation of HMF, indanone and guanidine produced the compound **91**, which was found to be a potent  $A_{2A}$  adenosine receptor antagonist.<sup>[107]</sup> A montmorillonite K-10 clay-catalyzed condensation of HMF, 2-aminopyridine and isocyanide was performed to generate the 1,8a-dihydroimidazo[1,2-*a*]pyridine derivative **92**.<sup>[108]</sup> More recently, our group disclosed the Biginelli reaction of 5-HMF using zinc chloride as Lewis acid catalyst, allowing access to furan-based 3,4-dihydropyrimidin-2-ones **93**.<sup>[109]</sup> The urea scope could be extended to thiourea and heterocyclic urea derivatives. Another interesting multicomponent reaction is the coupling of HMF, terminal aromatic alkynes, primary aliphatic amines and carbon dioxide using copper iodide as catalyst, to deliver oxazolidinones **94**.<sup>[110]</sup> In the case of anilines and secondary aliphatic amines, the  $A^3$  coupling (aldehyde, amine and alkyne) took place under same conditions providing compounds **95**.






## Scheme 36. MCR strategies involving 5-HMF


### 2.2.4 Oxidative transformations

The reaction of diazo derivatives **96** on 5-HMF catalyzed by *N*-heterocyclic carbene (NHC) allowed the preparation of acylhydrazones **97** displaying some antitumoral activity (Scheme 37).<sup>[111]</sup> An *in situ*-generated NHC catalyst (prepared by reacting imidazolium SiMes·HCl with DBU) was used to reverse the reactivity of the aldehyde group of HMF. The authors observed neither Cannizzaro reaction nor NHC-catalyzed benzoin condensation. In some cases, the expected acylhydrazones were obtained as mixtures of *E/Z* geometric isomers. The extension of the scope to hydroxymethyl protected substrates pointed out significant differences in terms of reactivity. For example, benzoyl HMF gave only 3% yield of acylhydrazone, while HMF benzyl and TBDMS ethers afforded 89% and 99% yields respectively.




Scheme 37. Reaction between 5-HMF and diazo derivatives

HMF has been converted to the nitrile **98** by treatment with an ammonia solution in the presence of molecular iodine in a mixture of water and THF (Scheme 38).<sup>[112]</sup> A similar oxidative cyanation was implemented in a one-pot procedure for converting HMF into amide **99**,<sup>[113]</sup> where the *in situ*-generated nitrile was hydrolyzed to the corresponding carboxamide under the action of a nitrile hydratase from *R. rhodochrous* IFO 15564. The same transformation could be performed under modified experimental conditions by flow process with hydroxylamine hydrochloride as nitrogen source.<sup>[114]</sup> The reaction was completed in 5 minutes taking advantage of the flow chemistry technique (Scheme 38).



Scheme 38. Conversion 5-HMF to nitrile **98** and amide **99**

Catalytic oxidation of 5-HMF in the presence of aqueous ammonia and manganese oxide octahedral molecular sieves (OMS-2) as catalyst afforded the bis-imidate **100** in good yield (Scheme 39). Subsequent treatment with ammonium chloride provided the corresponding bis-amidine **101**.<sup>[115]</sup>




Scheme 39. Transformation of 5-HMF to dicarboximidate and diamidine salt

## 2.3 Carbon-oxygen and carbon-sulfur bond formation reactions


The nucleophilic attack of alcohols on aldehyde groups forms acetals, which are often applied as strategies for carbonyl compounds protection due to the reversibility of the acetal formation. Treatment of HMF with trimethyl orthoformate in the presence of ytterbium sulfate supported on Amberlite 15 gave access to the acetal **102** in 80% yield (Scheme 40).<sup>[116]</sup> The same protected HMF was obtained in high yield by Al-beta-Zeolite-catalyzed acetalization in methanol.<sup>[117]</sup> Using cerium phosphate as a catalyst in alcohols, both cyclic and acyclic acetals have been obtained in good yields.<sup>[118]</sup> Amberlyst 15 was also found to be an efficient catalyst for the preparation

of cyclic acetals.<sup>[119]</sup>




**Scheme 40.** Acetalization of HMF with simple alcohols

The acetalization of HMF with glycerol was performed by means of solid acid catalysis, yielding a mixture of 1,3-dioxolane **105** and 1,3-dioxane **106** (Scheme 41).<sup>[120-121]</sup> The acetalization with glycerol was extended to several *in situ* formed 5-(alkyloxymethyl)furfurals for the preparation of potential surfactant molecules **107** and **108** in a one-pot process.<sup>[122]</sup> The resulting compounds with HLB values in the range of 4.9 to 6.6 and good thermal and water stability could be useful for industrial applications as W/O emulsifiers.



**Scheme 41.** The acetalization of HMF and its derivative with glycerol

Likewise, thioacetalization of HMF under various catalytic conditions has been also reported. The reaction with ethyl-2-mercaptoacetate in the presence of acidic H-β zeolite and 1-butyl-3-methylimidazolium chloride provided the corresponding dithioacetal **109** in 85% yield (Scheme 42).<sup>[123]</sup> Ou prepared dithioacetal **110** by the reaction of HMF and cysteine in order to evaluate its cytotoxicity because this dithioacetal could be formed during food processing and digestion.<sup>[124]</sup>



**Scheme 42.** Thioacetalization of 5-HMF

### 3 Reactions on the hydroxyl group


This section covers the strategies taking advantage of the reactivity of this benzylic alcohol to produce valuable chemicals. Apart from reactions where the hydroxymethyl group of HMF has been exploited for grafting it onto solid supports or polymers, several other transformations involving the CH<sub>2</sub>OH group of HMF were reported, such as conversion to esters, ethers, halides and amines, as well as its reactivity as an alkylating reagent in Friedel-Crafts alkylations.

#### 3.1 Conversion to halides

HMF can be easily converted to 5-chloromethylfurfural **111** or 5-bromomethylfurfural **112** *via* halogen substitution of the hydroxyl group. A range of classical halogenating conditions could be successfully applied to HMF (Table 4). The resulting halogenated HMF derivatives proved to be useful intermediates for the synthesis


of several valuable furan-based complex compounds. 5-Chloromethylfurfural **111** being easily and directly obtained from sugars is now also claimed as a platform chemical itself.<sup>[125]</sup>

**Table 4.** Halogenation experimental conditions



Entry	Product	Conditions	Yield	Ref.
1	<b>111</b>	HCl (gas), Et <sub>2</sub> O, 9 h	87%	[126]
2		HCl (aq), DCM or DCE, rt, 24 h	86-92%	[1, 127-128]
3		SOCl <sub>2</sub> , pyridine, DCE, 5 h	71%	[126]
4		Me <sub>3</sub> SiCl, CDCl <sub>3</sub> , 6 h	92%	[126]
5		POCl <sub>3</sub> , DMF, 5 h	77%	[129]
6	<b>112</b>	HBr (gas), Et <sub>2</sub> O, 5 h	64%	[126, 130]
7		HBr (aq), DCE, 65 °C, 1 h	94%	[127]
8		SOBr <sub>2</sub> , pyridine, DCE, 5 h	75%	[126]
9		PBr <sub>3</sub> , CaCO <sub>3</sub> , DCE, 5 h	62%	[126]
10		Me <sub>3</sub> SiBr, CHCl <sub>3</sub> or CHCl <sub>2</sub> CH <sub>2</sub> Cl, 3 h	>98%	[126, 131]
11		NBS, PPh <sub>3</sub> , DCM, -5 °C, 0.5 h	87%	[132]


Taking advantage of chlorination of HMF as a key step, Ananikov reported an efficient synthetic route to ranitidine, a medication which decreases stomach acid production (Scheme 43).<sup>[1]</sup> The synthetic route consisted of the nucleophilic substitution of 5-chloromethylfurfural **111** by *N*-acetylcysteamine. Subsequent transformations of the thioether **113** afforded smoothly ranitidine in 65% overall yield from HMF. Changing the reagent to an arylisocyanate in the last step provided CAP-1, a novel compound with inhibition of HIV-1 capsid protein assembly activity.<sup>[128]</sup>



**Scheme 43.** Synthesis of ranitidine and CAP-1 from 5-HMF

### 3.2 Conversion to ethers and acetals

HMF can undergo self-etherification to yield 5,5'-[oxybis(methylene)]bis-2-furfural (OBMF) **114** under acidic medium (Scheme 44). It is one of the (undesired) processes occurring during degradation of HMF upon storage.<sup>[1]</sup> The dehydration of HMF in the presence of *p*-TSA and phosphorus pentoxide in toluene with a continuous water removal using a Dean-Stark trap provides OBMF in 76% yield.<sup>[133]</sup> OBMF can also be obtained in 55% yield by thermal dehydration in DMSO without any acidic catalyst.<sup>[134]</sup> Additionally, several heterogeneous solid acids were reported to catalyze this reaction (e.g., Amberlyst-15, Amberlite IR 120, Preyssler heteropolyacids, graphene oxide).<sup>[1, 135-140]</sup> Finally, OBMF could also be obtained by Williamson reaction of HMF and 5-chloromethylfurfural.<sup>[141]</sup>



**Self-etherification**


Conditions A: *p*-TSA, P<sub>2</sub>O<sub>5</sub>, toluene, reflux, 76%  
 Conditions B: DMSO, 155 °C, 55%  
 Conditions C: Amberlyst-15, benzene, reflux, 79%  
 Conditions D: Amberlite IR 120 (H), benzene, reflux, 38%  
 Conditions E: Preyssler HPA, 70 °C, 84%  
 Conditions F: graphene oxide, DCM, 100 °C, 86%  
 Conditions G: Al-MCM-41, trifluorotoluene, 100 °C, 99%  
 Conditions H: Sn-Mont catalyst, DCM, 100 °C, 96%  
 Conditions I: Amberlyst-15, DCM, ultrasonication, 96%

**Williamson reaction**

Conditions: K<sub>2</sub>CO<sub>3</sub>, KI, acetone, reflux, 43%


**Scheme 44.** Conversion HMF to OBMF

Simple alkyl ethers (e.g. 5-ethoxymethylfurfural) can be prepared by catalytic etherification of HMF and alcohols at high temperature.<sup>[142-143]</sup> A milder alternative is the Williamson reaction between HMF and an alkyl halide. It was reported that treatment of HMF with sodium hydride/iodomethane could give the methyl ether in 94% yield.<sup>[22]</sup> HMF reacted with benzyl bromide in the presence of Ag<sub>2</sub>O in DMF to afford 5-benzyloxymethylfurfural in 72% yield.<sup>[135]</sup> Pichiafuran C, a monofuran metabolite isolated from *Pichia membranifaciens* yeast, was synthesized *via* Yb(OTf)<sub>3</sub>-catalyzed reaction of HMF with 2-phenylethanol and subsequent reduction of **115** with NaBH<sub>4</sub> (Scheme 45).<sup>[144]</sup>




**Scheme 45.** Synthesis of Pichiafuran C

Sessiline, an alkaloid isolated from the fruits of *Acanthopanax sessiliflorus* was prepared in 54% yield in neat conditions by reaction of 5-HMF with amidocarbinol **116** (obtained in 3 steps and 28% yield from succinimide).<sup>[145]</sup> The proposed mechanism is the nucleophilic addition of HMF onto the iminium intermediate, which is in equilibrium with the amidocarbinol (Scheme 46).



**Scheme 46.** Synthesis of Sessiline


Stevens and coworkers performed the β-alkoxylation of the *in situ* formed 3-imidoallenylphosphonate **117** with a range of alcohols (Scheme 47).<sup>[146]</sup> Despite a crude yield estimated to 90%, the product **118** was isolated only in 33% yield, probably due to partial degradation during reverse phase flash chromatography purification.



**Scheme 47.** Alkoxylation of 3-imidoallenylphosphonate with 5-HMF

Descotes prepared the tetrahydropyranyl ether **119** in 72% yield by reaction between HMF and dihydropyran catalyzed by pyridinium *p*-toluenesulfonate (Scheme 48).<sup>[147]</sup> The Ferrier rearrangement of HMF and

peracetylated glycal **120** under Lewis acid catalysis was reported by the same group in 2001 (Scheme 48).<sup>[148]</sup> The best yield was obtained when the combination LiBF<sub>4</sub>/SnCl<sub>2</sub> was employed as Lewis acid catalysts. More recently, Zhang described the transformation of various glycals including **120** with HMF in the presence of FeCl<sub>3</sub>·6H<sub>2</sub>O/C as a recyclable catalyst in DCE, leading to a range of 2,3-unsaturated glycosides in 67-92% yields.<sup>[149]</sup> Other carbohydrate derivatives of HMF have also been studied. Cottier investigated several possible pathways to prepare β-glucosylmethylfurfural **123** starting from HMF and glycosyl donors such as **122**.<sup>[116]</sup> The Ag<sub>2</sub>O-catalyzed coupling of HMF and glycosyl bromide **122a** gave only 11% yield of the β-glucosylmethylfurfural. The yield was improved to 32% using the imidate **122b** and boron trifluoride diethyl etherate (Scheme 48).




**Scheme 48.** Acetal formation

### 3.3 Esterifications


Classical esterification conditions were applied for the functionalization of the hydroxyl group of 5-HMF. The reaction with acetic anhydride in the presence of a base such as pyridine or sodium acetate, furnished 5-acetoxymethylfurfural in up to 93% yield.<sup>[16, 135]</sup> Benzoate esters can be prepared by reaction with benzoyl halides under similar conditions.<sup>[150-151]</sup>

The esterification of 4-*O*-acetylferulic acid with HMF was performed in pyridine using DCC as coupling reagent in the presence of *p*-toluenesulfonic acid, providing the corresponding ester **124** in 40% yield (Scheme 49).<sup>[152]</sup> The ester **125** was also prepared *via* DCC-promoted coupling reaction between HMF and the coumarin-derived acid, and its inhibitory activity against protein tyrosine phosphatase 1B was studied.<sup>[153]</sup>



**Scheme 49.** DCC-promoted esterification between HMF and carboxylic acids


Tamariz's group prepared the natural product **127**, isolated from the Noni fruit, by two successive esterifications: first, the esterification of HMF hydroxyl group with succinic anhydride leading to the intermediate ester-acid **126**, then the alkylation of this latter with *n*-butyl bromide (Scheme 50) leading to the diester **127**. The stepwise and one-pot process gave same overall yields (85%).<sup>[154]</sup>



**Scheme 50.** Synthesis of natural succinate derivative

Mumefural, a citrate ester of 5-HMF known to improve human blood fluidity, can be prepared from malic acid


and HMF.<sup>[155]</sup> The intermediate **128** (prepared from malic acid) reacted with HMF in the presence of DCC and a catalytic amount of DMAP in dichloromethane to deliver the ester **129** in 91% yield (Scheme 51). Subsequent deprotection gave access to mumeifural in 68% yield.



**Scheme 51.** Synthesis of mumeifural from HMF

Biocatalytic esterification of HMF with different acyl donors such as carboxylic acids and methyl or ethyl esters has been reported (Scheme 52).<sup>[156]</sup> Good to excellent yields of esters **130** were obtained using the lipase CAL-B as catalyst in solvent-free conditions. In the case of short-chain acids, a mixture of carboxylic acid and *tert*-butanol (1:1 v/v) was used instead of solvent-free conditions to avoid the deactivation of the lipase due to the high acidity of the medium. Interestingly the same enzyme proved also to be efficient to prepare methyl carbonate **131** from dimethyl carbonate as well as HMF-esters directly from soybean oil.

The enzymatic esterification between HMF and levulinic acid to levulinic acid **132** was reported by Li et al.<sup>[157]</sup> Novozym 435 was selected as the best catalyst and the reaction could proceed in alcoholic solvents such as *tert*-butanol, or 2-methyl-2-butanol as well as in ethereal solvents (CPME, or 2-MeTHF), with more than 90% conversion.




**Scheme 52.** Biocatalytic esterifications

### 3.4 Amination reactions


Strictly speaking, there are no examples of real amination on the OH group of HMF to make amino-deoxy-HMF. However, a few studies about the preparation of amides, hydrazines and ureas have been reported.

The group of Wierenga reported the direct preparation of hydrazine **133** under Mitsunobu conditions from 5-HMF.<sup>[158]</sup> In the absence of any other nucleophile, the reduced DEAD reagent acts as a nucleophile in SN2 reaction to produce compound **133** in 12% yield (Scheme 53). Comparatively, furfuryl alcohol or the acetal-protected HMF provided much higher yields (60% and 86% respectively) under the same conditions.




**Scheme 53.** Reaction between HMF and diethyl azodicarboxylate

Lewis et al. prepared the *N*-hydroxyurea derivative **134** by reacting HMF with *N,O*-(bisphenoxy-carbonyl)hydroxylamine under Mitsunobu conditions, followed by treatment with ammonia. Subsequent reductive amination with piperazine **135** released the final product **136** (yields not reported), which exhibited antihistaminergic activity (Scheme 54).<sup>[159]</sup>



**Scheme 54.** Reaction between HMF and *N,O*-(bisphenoxycarbonyl) hydroxylamine


The Ritter reaction was also applied to HMF, which gave access to the corresponding amides **137** in around 50% yield by treatment with trifluoromethanesulfonic acid in acetonitrile or benzonitrile at room temperature (Scheme 55).<sup>[135]</sup>



**Scheme 55.** Reaction between HMF and nitriles

### 3.5 Construction of a C-C(Ar) bonds

The Friedel–Crafts alkylation of arenes using HMF as the alkylating agent has been reported by different research groups. Beller et al. implemented the alkylation of *o*-xylene using  $\text{FeCl}_3$  (10 mol%) as a catalyst providing a mixture of regioisomers in 37% yield (ratio **138/139** of 62:38) (Scheme 56).<sup>[160]</sup> The reaction with mesitylene, either promoted by  $\text{FeCl}_3$  or nitrosonium tetrafluoroborate, provided the product **140** in excellent yield.<sup>[161-162]</sup> Heating HMF in refluxing toluene in the presence of poly(3,4-ethylenedioxythiophene) salt gave rise to a mixture of two isomers **141** and **142** in a global 81% yield.<sup>[163]</sup> Employing  $\text{Glu-Fe}_3\text{O}_4\text{-SO}_3\text{H}$  as a magnetic carbonaceous solid acid catalyst led to a regioisomeric mixture of **141** and **142** containing some traces of the *meta* isomer.<sup>[164]</sup> Recently, Vasilyev and coworkers reported the arylation of HMF with arenes in trifluoromethanesulfonic acid medium at room temperature.<sup>[165]</sup> 5-Arylmethylfurfurals **143** were prepared in 17–91% yields, along with by-products 2-arylmethyl-5-(diarylmethyl)furans **144** (5–37% yield) when highly nucleophilic arenes were used. Increasing the reaction time significantly affected the yield of the by-product. The same reactions using acidic zeolites CBV-720 as catalysts at 130 °C resulted in the selective formation of 5-arylmethylfurfurals (11–79% yield).




**Scheme 56.** Friedel–Crafts alkylation of arenes

## 4 Reactions on the furan ring

### 4.1 Ring opening/rearrangement


It is well known that HMF decomposes under acidic conditions at elevated temperature leading to the ring-opening product levulinic acid.<sup>[2,166]</sup> However, other derivatives arise from rearrangements of the furanic systems of HMF. For example, subjecting HMF to hydrothermolysis at high temperature and pressure led to 1,2,4-benzenetriol **145** as the major product in yields up to 46% with 50% HMF conversion (Scheme 57).<sup>[167]</sup> The other example is the conversion of HMF to phenols which was reported by Hashmi et al.<sup>[168]</sup> In this reaction,

HMF first undergoes a *O*-propargylation and the resulting aldehyde **146** is converted into trifurylmethane **147** under gold(III) catalysis. Finally, gold (I) catalysis promoted the conversion of **147** into the fused phenol **148** through ring rearrangement.




**Scheme 57.** Transformation of HMF and its derivatives to phenols

The hydrogenative/hydrolytic ring-opening of HMF in water provided 2,5-hexanedione (HD)<sup>[169-170]</sup> or 1-hydroxy-2,5-hexanedione (HHD)<sup>[171-172]</sup> depending on the catalytic conditions employed (Scheme 58). The latter was reported to be used as 1,4-diketone partner in the Paal-Knorr pyrrole synthesis.<sup>[173]</sup> HD and HHD can be converted into cyclopentanic derivatives, respectively 3-methyl-2-cyclopent-2-en-1-one (MCO)<sup>[174]</sup> and 2-hydroxy-3-methylcyclopent-2-enone (MCP)<sup>[175-176]</sup> *via* base-catalyzed intramolecular aldol reactions. Following an acid-catalyzed aldol reaction and subsequent hydrogenation sequence, HHD can be transformed into 3-hydroxymethyl cyclopentanone (HCPN)<sup>[177-178]</sup>. An alternative route toward cyclopentanic derivatives from HMF relies on a first catalytic reduction of the aldehyde generating 2,5-dihydroxymethylfuran (DHMF), followed by a Piancatelli rearrangement to 4-hydroxy-4-(hydroxymethyl)cyclopent-2-en-1-one (HHCPN) and further reduction of the produced cyclopentenone under the reaction conditions (Scheme 58).<sup>[179]</sup>



**Scheme 58.** Ring-opening/rearrangement of HMF derivatives

Oxidative conditions have also been employed to achieve ring opening transformations of intermediates arising from 5-HMF. Lichtenthaler and coworkers reported the preparation of various heterocycles such as the pyridazine **152**, the thiophene **153** or the pyrrole **154** from HMF (Scheme 59),<sup>[180]</sup> relying on a first reduction of the aldehyde and benzylation of the diol intermediate to produce **149**. This symmetrical disubstituted furan can undergo oxidative ring opening upon treatment with *m*-CPBA to yield diketone **150**. Chemoselective hydrogenation of **150** would provide diketone **151**. These 1,4-diketones were rapidly converted in heterocyclic scaffolds **152-154** *via* diverse transformations.






**Scheme 59.** Heterocycles obtained in three steps from HMF involving a furan ring-opening reactions

#### 4.2 Singlet oxygen oxidation

HMF can react with singlet oxygen *via* [4 + 2] cycloaddition to give rise to 5-hydroxy-5-(hydroxymethyl)-furan-2(5*H*)-one (**156**).<sup>[135]</sup> The cycloadduct **155** rearranged *via* a solvent-assisted decarbonylation reaction to produce efficiently the butenolide **156**. The reaction was recently reinvestigated by Kappe under continuous flow conditions (Scheme 60).<sup>[181]</sup> The best yield obtained was 93% using 1 mM rose bengal as photosensitizer in a mixture of *i*-PrOH/H<sub>2</sub>O as solvent with 0.5 mL/min flow rate.




**Scheme 60.** Reaction of HMF with singlet oxygen

#### 4.3 Diels-Alder reactions


Electron-rich furans have been widely employed as dienes in Diels-Alder reactions with electron-deficient dienophiles. In contrast to other furanic compounds, the presence of the strongly electron withdrawing aldehyde on the furan ring of HMF tends to deactivate the aromatic ring toward Diels-Alder reactions. In order to use HMF in this cycloaddition process, it usually requires chemical modification such as reduction toward 2,5-dihydroxymethylfuran (DHMF) or 2,5-dimethylfuran (DMF).

Ananikov and co-workers reported the synthesis of a series of oxabicyclic compounds starting from HMF and maleimide *via* one-pot reduction/cycloaddition/hydrogenation sequence or in a stepwise process taking advantage of a Diels-Alder reaction as the key step (Scheme 61).<sup>[182]</sup> Reduction of HMF to 2,5-dihydroxymethylfuran followed by *endo*-selective Diels-Alder cycloaddition with maleimide and hydrogenation in a single pot provided the product **157** in 73% yield (that could also be prepared in sequential-step procedure). The authors also applied the Diels-Alder reaction on HMF-derived bis(aminomethyl)furans, and bis(alkoxymethyl)furans and maleic anhydride could be employed as dienophile to prepare analogues of the anticancer drug norcantharidin.<sup>[183]</sup>



**Scheme 61.** Diels-Alder reaction of DHMF, bis(aminomethyl)furans and bis(alkoxymethyl)furans

Sheppard's group reported a strategy to efficiently react HMF with dienophiles, without changing the oxidation state of the aldehyde (Scheme 62).<sup>[184]</sup> HMF was first reacted with *N*-methylhydrazine to produce the corresponding hydrazone, before reacting with maleimides providing the cycloadducts **158**. The latter underwent deoxyaromatization to produce the phthalimides **159**. The reaction was carried out in water and the products were isolated in high yields by simple filtration.



### Scheme 62. Diels-Alder reaction of HMF-derived hydrazone


In addition, a range of HMF-derived furans such as 2,5-dimethylfuran (DMF) and dimethyl furan-2,5-dicarboxylate have been utilized as dienes in reactions with various dienophiles (e.g., ethylene, acrolein and benzyne) to accomplish the Diels-Alder reaction. Subsequent deoxygenation of the Diels-Alder adducts led to *p*-xylene or other substituted aromatic compounds.<sup>[185-193]</sup>

## 5 Multi-step and multi-functionalization sequences

As it has been shown throughout this review article, several methodological studies on the chemical modifications of HMF have been conducted, showing the high versatility of this C-6 synthon. In a synthetic perspective, 5-HMF has also been used in more complex multistep sequences, targeting complex and/or natural scaffolds.


In a study on intramolecular Diels-Alder reactions (IMDA) of furan derivatives, Sternbach used the 5-HMF as a platform for the preparation of the highly substituted keto-sulfone **160** (Scheme 63).<sup>[194]</sup> This substrate is poorly reactive in IMDA cycloaddition, however the conversion of the ketone to a cyanohydrin group using TMSCN/ZnI<sub>2</sub> reactivates the furan ring, which then can undergo an IMDA reaction in good yield producing tricyclic derivative **161** and epimer **161'** though with moderate diastereoselectivity.

A supported intramolecular Diels-Alder reaction was also performed on the furan ring of HMF after its coupling with a resin-bound amine (Scheme 63).<sup>[195]</sup> The polycyclic compound **163** could be obtained from the intermediate compound **162** after IMDA reaction followed by cleavage from the resin.



Scheme 63. Intramolecular Diels-Alder reaction on furan ring


Schinker et al. completed the stereoselective synthesis of furano-epothilone D, analog of natural epothilone which is a powerful antitumoral molecule, using HMF as a building block (Scheme 64).<sup>[196]</sup> Bromination of the hydroxymethyl group was the first step of functionalization. A sequence of a Negishi-type coupling and a diastereoselective aldol reaction with a chiral ketone produced the secondary alcohol **164**. This advanced intermediate was then converted to the furanic analogs of epothilone D. Both furano-epothilone D and its diastereoisomer displayed only moderate effect on tubulin assembly.



Scheme 64. Synthesis of furano-epothilone D from HMF


Loh *et al.* reported a synthetic route to natural products aspergillides A and B which exhibit potent cytotoxic activities towards mouse lymphocytic leukemia cell (Scheme 65).<sup>[197]</sup> In this route, all the carbon atoms in target molecules come from biomass-derived platform chemicals such as levulinic acid and HMF. The key steps in this synthesis were a Noyori's asymmetric transfer hydrogenation to produce enantiopure alcohol **165** which was then subjected to an Achmatowicz rearrangement and subsequent reduction leading to pyranes **166** and **167**,

advanced intermediates in the preparation of the natural products aspergillides A and B.




**Scheme 65.** Synthetic route to aspergillides A and B

A four-step process to prepare 8-oxa-3-aza-bicyclo[3.2.1]octane hydrochloride **169** starting from 5-HMF was described by Connolly et al.<sup>[198]</sup> Reduction of 5-HMF to the corresponding saturated diol followed by double tosylation and double  $S_N2$  with benzylamine generated the bicyclic intermediate **168**. The latter was hydrogenolyzed using Pearlman's catalyst in good yield to provide the final tricyclic adduct **169** (Scheme 66).




**Scheme 66.** Synthesis of 8-oxa-3-aza-bicyclo[3.2.1]octane hydrochloride **169**

MPEG-grafted 5-HMF **170** has been used as a substrate to prepare a series of tricyclic molecules **173** (Scheme 67).<sup>[199]</sup> The authors employed a 4-component Ugi reaction of supported HMF **170**, benzyl isocyanide and various primary amines and fumaric acid mono-amides (**171**) to generate intermediates **172**, that can undergo a tandem intramolecular Diels-Alder reaction of the furan ring followed by acidic workup to release final products **173**.




**Scheme 67.** Ugi/Diels-Alder tandem reaction

As illustrated above in Ugi/Diels-Alder reactions, the hydroxymethyl group present in 5-HMF offers a convenient handle for connection onto a solid support. The solid-supported HMF with a free carbonyl group could serve as a useful scaffold in combinatorial synthesis of furan-based libraries. Kundu and coworkers loaded HMF onto Sieber amide resin *via* the formation of amide and ester bonds using aliphatic dicarboxylic acid anhydride (Scheme 68).<sup>[200]</sup> The resin-supported HMFs **174** could be engaged in multicomponent  $A^3$  couplings to provide, after separation from the solid support, propargylamines **175**. Other multi-step transformations giving access to diverse nitrogen-containing heterocycles such as **176** and **177** were also successfully implemented using this supported synthesis strategy.




**Scheme 68.** Solid-phase combinatorial synthesis of HMF-based scaffolds

In 1956, Elming reported the synthesis of 6-hydroxymethyl-3-pyridinol **178** starting with reductive amination of HMF, followed by acetylation and electrolysis (Scheme 69).<sup>[87]</sup> The 3-pyridinol **178** could also be prepared by exposing 5-hydroxymethylfurfurylamine **179** to bromine in water/methanol at 0 °C.<sup>[201]</sup> Enantiopure synthesis of pyridinium salt (*S*)-alapyridaine **181** (a sweetness enhancer) was achieved under Br<sub>2</sub>/MeOH/H<sub>2</sub>O conditions from the amine formed by reductive amination of HMF with L-alanine.<sup>[202]</sup> Direct treatment of HMF with alkylamines in refluxing EtOH/H<sub>2</sub>O under basic conditions (pH adjusted ~9.4 with NaOH) has been a frequently-used method to prepared pyridinium inner salts **180** in the past.<sup>[203-205]</sup> Several pyridinium betaines derived from glycine, β-alanine and γ-aminobutyric acid proved to be potential bitter-suppressing candidates. More recently, an alternative method employing formic acid as a catalyst in EtOH/H<sub>2</sub>O to synthesize pyridinium inner salts **180** from HMF was reported by Afonso.<sup>[206]</sup>




**Scheme 69.** Conversion of 5-HMF to pyridinol and pyridinium salt

The preparation of key intermediates of synthetic routes towards 21,23-dioxaporphyrins **185** from HMF was reported as an alternative to known but complex synthetic procedures, addressing drawbacks such as low yields and purification issues (Scheme 70).<sup>[207]</sup> The addition of a phenyl-Grignard reagent on HMF, or the Henry reaction with nitromethane gave access to the furylcarbinol **182** and nitroalkene **183** respectively in good yields. Those intermediates were then reacted with pyrrole under indium chloride catalysis to generate diversely substituted bis-heteroarylmethanes **184**, highly valuable building-blocks for the preparation of 21,23-dioxaporphyrins **185**.



**Scheme 70.** Preparation of key intermediates to 21,23-dioxaporphyrin from 5-HMF

An example illustrating the usefulness of the olefination of HMF is the synthesis of trioxadispiroketal reported by Stockman.<sup>[208]</sup> In this strategy, HMF was subjected to tandem oxidation/Wittig reaction providing the diester **186** as a 6.6:1 mixture of (*E,E*)- and (*E,Z*)-isomers (Scheme 71). Reduction of both esters followed by hydrogenation of the two carbon-carbon double bonds on side chains afforded the corresponding diol **187**. Further treatment with *N*-bromosuccinimide furnished the [5,5,5]-trioxadispiroketal **188**. [6,5,6]-Trioxadispiroketal **189** was also prepared *via* a similar strategy using a homologated phosphonium ylide.



## Scheme 71. Application of the olefination of HMF to the synthesis of trioxadispiroketal

### 6 Conclusion

As it has been shown throughout this review article, 5-HMF has been the object of numerous studies on its use in fine chemical synthesis. Its relatively modest stability has been shown to be overcome by choosing adequately mild reaction conditions, notably with respect to acid-catalyzed reactions. Thanks to the presence of different functional groups on this platform molecule, it proved to be an excellent starting substrate for the preparation of analogs of drugs, photo-responsive molecules or complex heterocyclic scaffolds. The use of this C-6 synthon in novel synthetic routes is appealing, as it allows the incorporation of renewable carbon-sources into the final targets. The commercial availability of HMF being now fully established, it is expected to exhibit, in a foreseeable future, a growing number of such applications toward complex chemicals.

### 7 Acknowledgements

Financial support from CNRS, UCBL, INSA Lyon and CPE Lyon is gratefully acknowledged, as well as the ANR for a post-doctoral fellowship to C. V. (FurCab project, ANR-15-CE07-0016) and the China Scholarship Council for PhD grants to W. F.

### 8 References

- [1] Galkin, K. I.; Krivodaeva, E. A.; Romashov, L. V.; Zalesskiy, S. S.; Kachala, V. V.; Burykina, J. V.; Ananikov, V. P. Critical Influence of 5-Hydroxymethylfurfural Aging and Decomposition on the Utility of Biomass Conversion in Organic Synthesis. *Angew. Chem. Int. Ed.*, **2016**, *55* (29), 8338-8342.
- [2] Girisuta, B.; Janssen, L. P. B. M.; Heeres, H. J. A kinetic study on the decomposition of 5-hydroxymethylfurfural into levulinic acid. *Green Chem.*, **2006**, *8* (8), 701-709.
- [3] Nikolov, P. Y.; Yaylayan, V. A. Thermal Decomposition of 5-(Hydroxymethyl)-2-furaldehyde (HMF) and Its Further Transformations in the Presence of Glycine. *J. Agric. Food Chem.*, **2011**, *59* (18), 10104-10113.
- [4] Gomes, R. F. A.; Mitrev, Y. N.; Simeonov, S. P.; Afonso, C. A. M. Going Beyond the Limits of the Biorenewable Platform: Sodium Dithionite-Promoted Stabilization of 5-Hydroxymethylfurfural. *ChemSusChem*, **2018**, *11* (10), 1612-1616.
- [5] Rosatella, A. A.; Simeonov, S. P.; Frade, R. F. M.; Afonso, C. A. M. 5-Hydroxymethylfurfural (HMF) as a building block platform: Biological properties, synthesis and synthetic applications. *Green Chem.*, **2011**, *13* (4), 754-793.
- [6] de Vries, J. G. Chapter Eight - Green Syntheses of Heterocycles of Industrial Importance. 5-Hydroxymethylfurfural as a Platform Chemical. In *Adv. Heterocycl. Chem.*, Scriven, E. F. V.; Ramsden, C. A., Eds. Academic Press: **2017**; Vol. *121*, pp 247-293.
- [7] van Putten, R.-J.; van der Waal, J. C.; de Jong, E.; Rasrendra, C. B.; Heeres, H. J.; de Vries, J. G. Hydroxymethylfurfural, A Versatile Platform Chemical Made from Renewable Resources. *Chem. Rev.*, **2013**, *113* (3), 1499-1597.
- [8] Kucherov, F. A.; Romashov, L. V.; Galkin, K. I.; Ananikov, V. P. Chemical Transformations of Biomass-Derived C6-Furanic Platform Chemicals for Sustainable Energy Research, Materials Science, and Synthetic Building Blocks. *ACS Sustainable Chem. Eng.*, **2018**, *6* (7), 8064-8092.
- [9] Liu, D.; Zhang, Y.; Chen, E. Y. X. Organocatalytic upgrading of the key biorefining building block by a catalytic ionic liquid and N-heterocyclic carbenes. *Green Chem.*, **2012**, *14* (10), 2738-2746.
- [10] Liu, D.; Chen, E. Y. X. Diesel and Alkane Fuels From Biomass by Organocatalysis and Metal-Acid Tandem Catalysis. *ChemSusChem*, **2013**, *6* (12), 2236-2239.
- [11] Liu, D.; Chen, E. Y. X. Integrated Catalytic Process for Biomass Conversion and Upgrading to C12 Furoin and Alkane Fuel. *ACS Catal.*, **2014**, *4* (5), 1302-1310.
- [12] Wang, L.; Chen, E. Y. X. Recyclable Supported Carbene Catalysts for High-Yielding Self-Condensation of Furaldehydes into C10 and C12 Furoins. *ACS Catal.*, **2015**, *5* (11), 6907-6917.
- [13] Zang, H.; Chen, E. Y. X. Organocatalytic Upgrading of Furfural and 5-Hydroxymethyl Furfural to C10 and C12 Furoins with Quantitative Yield and Atom-Efficiency. *Int. J. Mol. Sci.*, **2015**, *16* (4), 7143-7158.
- [14] Yan, B.; Zang, H.; Jiang, Y.; Yu, S.; Chen, E. Y. X. Recyclable montmorillonite-supported thiazolium

- ionic liquids for high-yielding and solvent-free upgrading of furfural and 5-hydroxymethylfurfural to C10 and C12 furoins. *RSC Adv.*, **2016**, 6 (80), 76707-76715.
- [15] Donnelly, J.; Muller, C. R.; Wiermans, L.; Chuck, C. J.; Dominguez de Maria, P. Upgrading biogenic furans: blended C10-C12 platform chemicals via lyase-catalyzed carboligations and formation of novel C12 - choline chloride-based deep-eutectic-solvents. *Green Chem.*, **2015**, 17 (5), 2714-2718.
- [16] Coelho, J. A. S.; Trindade, A. F.; Andre, V.; Teresa Duarte, M.; Veiros, L. F.; Afonso, C. A. M. Trienamines derived from 5-substituted furfurals: remote  $\epsilon$ -functionalization of 2,4-dienals. *Org. Biomol. Chem.*, **2014**, 12 (46), 9324-9328.
- [17] Gomes, R. F. A.; Coelho, J. A. S.; Frade, R. F. M.; Trindade, A. F.; Afonso, C. A. M. Synthesis of Symmetric Bis(N-alkylaniline)triarylmethanes via Friedel-Crafts-Catalyzed Reaction between Secondary Anilines and Aldehydes. *J. Org. Chem.*, **2015**, 80 (20), 10404-10411.
- [18] Yu, C.; Liu, B.; Hu, L. Efficient Baylis-Hillman Reaction Using Stoichiometric Base Catalyst and an Aqueous Medium. *J. Org. Chem.*, **2001**, 66 (16), 5413-5418.
- [19] Yu, C.; Hu, L. Successful Baylis-Hillman Reaction of Acrylamide with Aromatic Aldehydes. *J. Org. Chem.*, **2002**, 67 (1), 219-223.
- [20] Tan, J.-N.; Ahmar, M.; Queneau, Y. HMF derivatives as platform molecules: aqueous Baylis-Hillman reaction of glucosyloxymethyl-furfural towards new biobased acrylates. *RSC Adv.*, **2013**, 3 (39), 17649-17653.
- [21] Tan, J.-N.; Ahmar, M.; Queneau, Y. Bio-based solvents for the Baylis-Hillman reaction of HMF. *RSC Adv.*, **2015**, 5 (85), 69238-69242.
- [22] Quiroz-Florentino, H.; Aguilar, R.; Santoyo, B. M.; Díaz, F.; Tamariz, J. Total Syntheses of Natural Furan Derivatives Rehmanones A, B, and C. *Synthesis*, **2008**, 2008 (7), 1023-1028.
- [23] Pupovac, K.; Palkovits, R. Cu/MgAl<sub>2</sub>O<sub>4</sub> as Bifunctional Catalyst for Aldol Condensation of 5-Hydroxymethylfurfural and Selective Transfer Hydrogenation. *ChemSusChem*, **2013**, 6 (11), 2103-2110.
- [24] Lewis, J. D.; Van de Vyver, S.; Román-Leshkov, Y. Acid-Base Pairs in Lewis Acidic Zeolites Promote Direct Aldol Reactions by Soft Enolization. *Angew. Chem. Int. Ed.*, **2015**, 54 (34), 9835-9838.
- [25] Lee, R.; Vanderveen, J. R.; Champagne, P.; Jessop, P. G. CO<sub>2</sub>-Catalysed aldol condensation of 5-hydroxymethylfurfural and acetone to a jet fuel precursor. *Green Chem.*, **2016**, 18 (19), 5118-5121.
- [26] Yutthalekha, T.; Suttipat, D.; Salakhum, S.; Thivasasith, A.; Nokbin, S.; Limtrakul, J.; Wattanakit, C. Aldol condensation of biomass-derived platform molecules over amine-grafted hierarchical FAU-type zeolite nanosheets (Zeolean) featuring basic sites. *Chem. Commun.*, **2017**, 53 (90), 12185-12188.
- [27] Bohre, A.; Saha, B.; Abu-Omar, M. M. Catalytic Upgrading of 5-Hydroxymethylfurfural to Drop-in Biofuels by Solid Base and Bifunctional Metal-Acid Catalysts. *ChemSusChem*, **2015**, 8 (23), 4022-4029.
- [28] Skowronski, R.; Grabowski, G.; Lewkowski, J.; Descotes, G.; Cottier, L.; Neyret, C. New chemical conversions of 5-hydroxymethylfurfural and the electrochemical oxidation of its derivatives. *Org. Prep. Proced. Int.*, **1993**, 25 (3), 353-355.
- [29] Arias, K. S.; Climent, M. J.; Corma, A.; Iborra, S. Chemicals from Biomass: Synthesis of Biologically Active Furanochalcones by Claisen-Schmidt Condensation of Biomass-Derived 5-hydroxymethylfurfural (HMF) with Acetophenones. *Top. Catal.*, **2016**, 59 (13), 1257-1265.
- [30] Zhao, F.; Zhao, Q.-J.; Zhao, J.-X.; Zhang, D.-Z.; Wu, Q.-Y.; Jin, Y.-S. Synthesis and cdc25B inhibitory activity evaluation of chalcones. *Chem. Nat. Compd.*, **2013**, 49 (2), 206-214.
- [31] Wang, Y.-H.; Dong, H.-H.; Zhao, F.; Wang, J.; Yan, F.; Jiang, Y.-Y.; Jin, Y.-S. The synthesis and synergistic antifungal effects of chalcones against drug resistant *Candida albicans*. *Bioorg. Med. Chem. Lett.*, **2016**, 26 (13), 3098-3102.
- [32] Suryawanshi, S. N.; Chandra, N.; Kumar, P.; Porwal, J.; Gupta, S. Chemotherapy of leishmaniasis part-VIII: Synthesis and bioevaluation of novel chalcones. *Eur. J. Med. Chem.*, **2008**, 43 (11), 2473-2478.
- [33] Mugunthan, G.; Ramakrishna, K.; Sriram, D.; Yogeewari, P.; Ravindranathan Kartha, K. P. Synthesis and screening of (E)-1-( $\beta$ -d-galactopyranosyl)-4-(aryl)but-3-ene-2-one against *Mycobacterium tuberculosis*. *Bioorg. Med. Chem. Lett.*, **2011**, 21 (13), 3947-3950.
- [34] Muthusamy, K.; Lalitha, K.; Prasad, Y. S.; Thamizhanban, A.; Sridharan, V.; Maheswari, C. U.; Nagarajan, S. Lipase-Catalyzed Synthesis of Furan-Based Oligoesters and their Self-Assembly-Assisted Polymerization. *ChemSusChem*, **2018**, 11 (14), 2453-2463.
- [35] Lockman, J. W.; Reeder, M. D.; Suzuki, K.; Ostanin, K.; Hoff, R.; Bhoite, L.; Austin, H.; Baichwal, V.; Adam Willardsen, J. Inhibition of eEF2-K by thieno[2,3-b]pyridine analogues. *Bioorg. Med. Chem. Lett.*, **2010**, 20 (7), 2283-2286.
- [36] Meguellati, A.; Ahmed-Belkacem, A.; Yi, W.; Haudecoeur, R.; Crouillère, M.; Brillet, R.; Pawlotsky, J.-M.; Boumendjel, A.; Peuchmaur, M. B-ring modified auronones as promising allosteric inhibitors of hepatitis C virus RNA-dependent RNA polymerase. *Eur. J. Med. Chem.*, **2014**, 80, 579-592.
- [37] Taylor, K. M.; Taylor, Z. E.; Handy, S. T. Rapid synthesis of auronones under mild conditions using a combination of microwaves and deep eutectic solvents. *Tetrahedron Lett.*, **2017**, 58 (3), 240-241.

- [38] Witczak, Z. J.; Bielski, R.; Mencer, D. E. Concise and efficient synthesis of *E*-stereoisomers of exocyclic carbohydrate enones. Aldol condensation of dihydrolevoglucosenone with five-membered aromatic aldehydes Part 1. *Tetrahedron Lett.*, **2017**, *58* (43), 4069-4072.
- [39] Zhao, F.; Dong, H.-H.; Wang, Y.-H.; Wang, T.-Y.; Yan, Z.-H.; Yan, F.; Zhang, D.-Z.; Cao, Y.-Y.; Jin, Y.-S. Synthesis and synergistic antifungal effects of monoketone derivatives of curcumin against fluconazole-resistant *Candida* spp. *MedChemComm*, **2017**, *8* (5), 1093-1102.
- [40] Liu, Q.; Zhang, C.; Shi, N.; Zhang, X.; Wang, C.; Ma, L. Production of renewable long-chained cycloalkanes from biomass-derived furfurals and cyclic ketones. *RSC Adv.*, **2018**, *8* (25), 13686-13696.
- [41] Amarasekara, A. S.; Singh, T. B.; Larkin, E.; Hasan, M. A.; Fan, H.-J. NaOH catalyzed condensation reactions between levulinic acid and biomass derived furan-aldehydes in water. *Ind. Crops Prod.*, **2015**, *65*, 546-549.
- [42] Keskinvälti, J.; Wrigstedt, P.; Lagerblom, K.; Repo, T. One-step Pd/C and Eu(OTf)<sub>3</sub> catalyzed hydrodeoxygenation of branched C11 and C12 biomass-based furans to the corresponding alkanes. *Appl. Catal. A: Gen.*, **2017**, *534*, 40-45.
- [43] Lin, L.; Shi, Q.; Nyarko, A. K.; Bastow, K. F.; Wu, C.-C.; Su, C.-Y.; Shih, C. C. Y.; Lee, K.-H. Antitumor Agents. 250. Design and Synthesis of New Curcumin Analogues as Potential Anti-Prostate Cancer Agents. *J. Med. Chem.*, **2006**, *49* (13), 3963-3972.
- [44] Martichonok, V. V.; Chiang, P. K.; Dornbush, P. J.; Land, K. M. On Regioselectivity of Aldol Condensation of Aromatic Aldehydes with Borate Complex of Acetylacetone. *Synth. Commun.*, **2014**, *44* (9), 1245-1250.
- [45] Shao, W.-Y.; Cao, Y.-N.; Yu, Z.-W.; Pan, W.-J.; Qiu, X.; Bu, X.-Z.; An, L.-K.; Huang, Z.-S.; Gu, L.-Q.; Chan, A. S. C. Facile preparation of new unsymmetrical curcumin derivatives by solid-phase synthesis strategy. *Tetrahedron Lett.*, **2006**, *47* (24), 4085-4089.
- [46] Tarleton, M.; Dyson, L.; Gilbert, J.; Sakoff, J. A.; McCluskey, A. Focused library development of 2-phenylacrylamides as broad spectrum cytotoxic agents. *Biorg. Med. Chem.*, **2013**, *21* (1), 333-347.
- [47] Tarleton, M.; Gilbert, J.; Sakoff, J. A.; McCluskey, A. Cytotoxic 2-phenylacrylnitriles, the importance of the cyanide moiety and discovery of potent broad spectrum cytotoxic agents. *Eur. J. Med. Chem.*, **2012**, *57*, 65-73.
- [48] Feng, Z.; Jia, J.; Liu, Y.; Wang, Z.; Zhao, X. 3-furyl-2-cyano-2-acrylamide derivative, preparation method therefor, pharmaceutical composition and use thereof US20160272604A1, 2016.
- [49] Hanefeld, W.; Schlitzer, M.; Debski, N.; Euler, H. 3-(2,5-Dioxopyrrolidin-1-yl), 3-(2,6-dioxopiperidin-1-yl), and 3-(1,3-dioxoisindolin-2-yl)rhodanines. a novel type of rhodanine derivatives. *J. Heterocycl. Chem.*, **1996**, *33* (4), 1143-1146.
- [50] Gregg, B. T.; Golden, K. C.; Quinn, J. F.; Tymoshenko, D. O.; Earley, W. G.; Maynard, D. A.; Razzano, D. A.; Rennells, W. M.; Butcher, J. Expedient Lewis Acid Catalyzed Synthesis of a 3-Substituted 5-Arylidene-1-methyl-2-thiohydantoin Library. *J. Comb. Chem.*, **2007**, *9* (6), 1036-1040.
- [51] Lukevics, E.; Ignatovich, L.; Shestakova, I. Synthesis, psychotropic and anticancer activity of 2,2-dimethyl-5-[5'-trialkylgermyl(silyl)-2'-hetarylidene]-1,3-dioxane-4,6-diones and their analogues. *Appl. Organomet. Chem.*, **2003**, *17* (12), 898-905.
- [52] Gomes, R. F. A.; Coelho, J. A. S.; Afonso, C. A. M. Direct Conversion of Activated 5-Hydroxymethylfurfural into delta-Lactone-Fused Cyclopentenones. *ChemSusChem*, **2019**, *12* (2), 420-425.
- [53] Nikolov, M. N.; Poneva, M. V. Spectral Determination of the Structure of 5-Hydroxymethylfurfurylidene Barbituric Acid. *Spectrosc. Lett.*, **1987**, *20* (10), 821-834.
- [54] Mercep, M.; Malnar, I.; Filipovic Susic, A.; Mesic, M. Preparation of antiinflammatory conjugates of erythromycin macrolides and coumarins. WO2006092739A1, 2006.
- [55] van Schijndel, J.; Canalle, L. A.; Molendijk, D.; Meuldijk, J. Exploration of the Role of Double Schiff Bases as Catalytic Intermediates in the Knoevenagel Reaction of Furanic Aldehydes: Mechanistic Considerations. *Synlett*, **2018**, *29* (15), 1983-1988.
- [56] Cummings, R. T.; DiZio, J. P.; Krafft, G. A. Photoactivable fluorophores. 2. Synthesis and photoactivation of functionalized 3-aryl-2-(2-furyl)-chromones. *Tetrahedron Lett.*, **1988**, *29* (1), 69-72.
- [57] Miller, R. Synthesis and Stereochemistry of (E)-5-(3,4,5,6-Tetrahydropyrid-3-ylidenemethyl)-2-furanmethanol, a Product of the Reaction between D-Glucose and L-Lysine. *Acta Chem. Scand. B*, **1987**, *41*, 208-209.
- [58] Shao, X.; Li, Z.; Qian, X.; Xu, X. Design, Synthesis, and Insecticidal Activities of Novel Analogues of Neonicotinoids: Replacement of Nitromethylene with Nitroconjugated System. *J. Agric. Food Chem.*, **2009**, *57* (3), 951-957.
- [59] Lee, J.-S.; Kang, N.-y.; Kim, Y. K.; Samanta, A.; Feng, S.; Kim, H. K.; Vendrell, M.; Park, J. H.; Chang, Y.-T. Synthesis of a BODIPY Library and Its Application to the Development of Live Cell Glucagon Imaging Probe. *J. Am. Chem. Soc.*, **2009**, *131* (29), 10077-10082.

- [60] Karaguni, I.-M.; Glüsenkamp, K.-H.; Langerak, A.; Geisen, C.; Ullrich, V.; Winde, G.; Möröy, T.; Müller, O. New indene-derivatives with anti-proliferative properties. *Bioorg. Med. Chem. Lett.*, **2002**, *12* (4), 709-713.
- [61] Mouloungui, Z.; Delmas, M.; Gaset, A. Synthesis of  $\alpha,\beta$  Unsaturated Esters Using a Solid-Liquid Phase Transfer in a Slightly Hydrated Aprotic Medium. *Synth. Commun.*, **1984**, *14* (8), 701-706.
- [62] Mouloungui, Z.; Delmas, M.; Gaset, A. Synthesis of  $\alpha, \beta$ -Ethylenic Esters in a Heterogenous Solid-Liquid Medium. II - A Transesterification Reaction Linked to a Wittig-Horner Reaction in a Protic Medium. *Synth. Commun.*, **1985**, *15* (6), 491-494.
- [63] Goodman, S. N.; Jacobsen, E. N. A Practical Synthesis of  $\alpha,\beta$ -Unsaturated Imides, Useful Substrates For Asymmetric Conjugate Addition Reactions. *Adv. Synth. Catal.*, **2002**, *344* (9), 953-956.
- [64] Almirante, N.; Cerri, A.; Fedrizzi, G.; Marazzi, G.; Santagostino, M. A general, [1+4] approach to the synthesis of 3(5)-substituted pyrazoles from aldehydes. *Tetrahedron Lett.*, **1998**, *39* (20), 3287-3290.
- [65] Fumagalli, T.; Sello, G.; Orsini, F. One-Pot, Fluoride-Promoted Wittig Reaction. *Synth. Commun.*, **2009**, *39* (12), 2178-2195.
- [66] Yoshida, N.; Kasuya, N.; Haga, N.; Fukuda, K. Brand-new Biomass-based Vinyl Polymers from 5-Hydroxymethylfurfural. *Polym. J.*, **2008**, *40*, 1164-1169.
- [67] Han, M.; Liu, X.; Zhang, X.; Pang, Y.; Xu, P.; Guo, J.; Liu, Y.; Zhang, S.; Ji, S. 5-Hydroxymethyl-2-vinylfuran: a biomass-based solvent-free adhesive. *Green Chem.*, **2017**, *19* (3), 722-728.
- [68] Romashov, L. V.; Ananikov, V. P. Alkynylation of Bio-Based 5-Hydroxymethylfurfural to Connect Biomass Processing with Conjugated Polymers and Furanic Pharmaceuticals. *Chem. Asian J.*, **2017**, *12* (20), 2652-2655.
- [69] Sharma, U. K.; Gemoets, H. P. L.; Schröder, F.; Noël, T.; Van der Eycken, E. V. Merger of Visible-Light Photoredox Catalysis and C-H Activation for the Room-Temperature C-2 Acylation of Indoles in Batch and Flow. *ACS Catal.*, **2017**, *7* (6), 3818-3823.
- [70] Ramonczai, J.; Vargha, L. Studies on Furan Compounds. III. A New Synthesis of Furyl Ketones. *J. Am. Chem. Soc.*, **1950**, *72* (6), 2737-2737.
- [71] Nicklaus, C. M.; Minnaard, A. J.; Feringa, B. L.; de Vries, J. G. Synthesis of renewable fine-chemical building blocks by reductive coupling between furfural derivatives and terpenes. *ChemSusChem*, **2013**, *6* (9), 1631-1635.
- [72] Hirapara, P.; Riemer, D.; Hazra, N.; Gajera, J.; Finger, M.; Das, S. CO<sub>2</sub>-assisted synthesis of non-symmetric  $\alpha$ -diketones directly from aldehydes via C-C bond formation. *Green Chem.*, **2017**, *19* (22), 5356-5360.
- [73] Cooper, W. F.; Nuttall, W. H. CXII.—Furan-2 : 5-dialdehyde. *J. Chem. Soc., Trans.*, **1912**, *101*, 1074-1081.
- [74] Middendorp, J. A. Sur l'oxyméthylfurfuro. *Recl. Trav. Chim. Pays-Bas*, **1919**, *38* (1), 1-71.
- [75] Cottier, L.; Descotes, G.; Lewkowski, J.; Skowroński, R. Synthesis and its stereochemistry of aminophosphonic acids derived from 5-hydroxymethylfurfural. *Phosphorus, Sulfur Silicon Relat. Elem.*, **1996**, *116* (1), 93-100.
- [76] Fan, W.; Queneau, Y.; Popowycz, F. The synthesis of HMF-based  $\alpha$ -amino phosphonates via one-pot Kabachnik-Fields reaction. *RSC Adv.*, **2018**, *8* (55), 31496-31501.
- [77] Amarasekara, A., S. ; Edigin, O.; Hernandez, W. Cycloaddition Reactions of 5-Hydroxymethyl-Furan-2-Nitrileoxide. *Lett. Org. Chem.*, **2007**, *4* (5), 306-308.
- [78] Wang, E.; Zhou, Y.; Huang, Q.; Pang, L.; Qiao, H.; Yu, F.; Gao, B.; Zhang, J.; Min, Y.; Ma, T. 5-Hydroxymethylfurfural modified rhodamine B dual-function derivative: Highly sensitive and selective optical detection of pH and Cu<sup>2+</sup>. *Spectrochim. Acta A*, **2016**, *152*, 327-335.
- [79] Alsaeedi, H. S.; Aljaber, N. A.; Ara, I. Synthesis and Investigation of Antimicrobial Activity of Some Nifuroxazide Analogues. *Asian J. Chem.*, **2015**, *27* (10), 3639-3646.
- [80] Brown, M. L.; Cheung, M.; Dickerson, S. H.; Drewry, D. H.; Lackey, K. E.; Peat, A. J.; Thomson, S. A.; Veal, J. M.; Wilson, J. L. R. Preparation of pyrazolopyrimidines as kinase inhibitors for the treatment of type 2 diabetes. WO2004009596A2, 2004.
- [81] Alturiqui, A. S.; Alaghaz, A.-N. M. A.; Ammar, R. A.; Zayed, M. E. Synthesis, Spectral Characterization, and Thermal and Cytotoxicity Studies of Cr(III), Ru(III), Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) Complexes of Schiff Base Derived from 5-Hydroxymethylfuran-2-carbaldehyde. *Journal of Chemistry*, **2018**, *2018*, 1-17.
- [82] Mohamed, O. G.; Khalil, Z. G.; Capon, R. J. Prolineimines: N-Amino-1-Pro-methyl Ester (Hydrazine) Schiff Bases from a Fish Gastrointestinal Tract-Derived Fungus, *Trichoderma* sp. CMB-F563. *Org. Lett.*, **2018**, *20* (2), 377-380.
- [83] Cisneros, L.; Serna, P.; Corma, A. Selective Reductive Coupling of Nitro Compounds with Aldehydes to Nitrones in H<sub>2</sub> Using Carbon-Supported and -Decorated Platinum Nanoparticles. *Angew. Chem. Int. Ed.*, **2014**, *53* (35), 9306-9310.



- [84] Pezzetta, C.; Veiros, L. F.; Oble, J.; Poli, G. Murai Reaction on Furfural Derivatives Enabled by Removable N,N'-Bidentate Directing Groups. *Chem. Eur. J.*, **2017**, *23* (35), 8385-8389.
- [85] Siopa, F.; Ramis Cladera, V.-A.; Afonso, C. A. M.; Oble, J.; Poli, G. Ruthenium-Catalyzed C-H Arylation and Alkenylation of Furfural Imines with Boronates. *Eur. J. Org. Chem.*, **2018**, *2018* (44), 6101-6106.
- [86] Cukalovic, A.; Stevens, C. V. Production of biobased HMF derivatives by reductive amination. *Green Chem.*, **2010**, *12* (7), 1201-1206.
- [87] Elming, N.; Clauson-Kaas, N. Transformation of 2-(hydroxymethyl)-5-(aminomethyl)-furan into 6-methyl-3-pyridinol. *Acta Chem. Scand.*, **1956**, *10*, 1603-1605.
- [88] Xu, Z.; Yan, P.; Xu, W.; Jia, S.; Xia, Z.; Chung, B.; Zhang, Z. C. Direct reductive amination of 5-hydroxymethylfurfural with primary/secondary amines via Ru-complex catalyzed hydrogenation. *RSC Adv.*, **2014**, *4* (103), 59083-59087.
- [89] Chatterjee, M.; Ishizaka, T.; Kawanami, H. Reductive amination of furfural to furfurylamine using aqueous ammonia solution and molecular hydrogen: an environmentally friendly approach. *Green Chem.*, **2016**, *18* (2), 487-496.
- [90] Komanoya, T.; Kinemura, T.; Kita, Y.; Kamata, K.; Hara, M. Electronic Effect of Ruthenium Nanoparticles on Efficient Reductive Amination of Carbonyl Compounds. *J. Am. Chem. Soc.*, **2017**, *139* (33), 11493-11499.
- [91] Chandra, D.; Inoue, Y.; Sasase, M.; Kitano, M.; Bhaumik, A.; Kamata, K.; Hosono, H.; Hara, M. A high performance catalyst of shape-specific ruthenium nanoparticles for production of primary amines by reductive amination of carbonyl compounds. *Chem. Sci.*, **2018**, *9* (27), 5949-5956.
- [92] Jagadeesh, R. V.; Murugesan, K.; Alshammari, A. S.; Neumann, H.; Pohl, M.-M.; Radnik, J.; Beller, M. MOF-derived cobalt nanoparticles catalyze a general synthesis of amines. *Science*, **2017**, *358* (6361), 326-332.
- [93] Zhu, M.-M.; Tao, L.; Zhang, Q.; Dong, J.; Liu, Y.-M.; He, H.-Y.; Cao, Y. Versatile CO-assisted direct reductive amination of 5-hydroxymethylfurfural catalyzed by a supported gold catalyst. *Green Chem.*, **2017**, *19* (16), 3880-3887.
- [94] Carrillo, A. I.; Llanes, P.; Pericàs, M. A. A versatile, immobilized gold catalyst for the reductive amination of aldehydes in batch and flow. *React. Chem. Eng.*, **2018**, *3* (5), 714-721.
- [95] Niphakis, M. J.; Gay, B. C.; Hong, K. H.; Bleeker, N. P.; Georg, G. I. Synthesis and evaluation of the anti-proliferative and NF- $\kappa$ B activities of a library of simplified tylophorine analogs. *Biorg. Med. Chem.*, **2012**, *20* (19), 5893-5900.
- [96] Kojiri, K.; Kondo, H.; Arakawa, H.; Ohkubo, M.; Suda, H. Preparation of indolopyrrolocarbazole derivatives having glucopyranosyl group and antitumor agents containing them. US6703373B1, 2004.
- [97] Pliitta, B.; Adamska, E.; Giel-Pietraszuk, M.; Fedoruk-Wyszomirska, A.; Naskręt-Barciszewska, M.; Markiewicz, W. T.; Barciszewski, J. New cytosine derivatives as inhibitors of DNA methylation. *Eur. J. Med. Chem.*, **2012**, *55*, 243-254.
- [98] Xu, Z.; Yan, P.; Liu, K.; Wan, L.; Xu, W.; Li, H.; Liu, X.; Zhang, Z. C. Synthesis of Bis(hydroxymethylfurfuryl)amine Monomers from 5-Hydroxymethylfurfural. *ChemSusChem*, **2016**, *9* (11), 1255-1258.
- [99] Sattler, L.; Zerban, F. W.; Clark, G. L.; Chu, C.-C. The Reaction of 2-Aminobenzenethiol with Al-doses and with Hydroxymethylfurfural. *J. Am. Chem. Soc.*, **1951**, *73* (12), 5908-5910.
- [100] Koehler, A. N.; Stefan, E.; Caballero, F. Preparation of fused 1,3-azole derivatives useful for the treatment of proliferative diseases. WO2016094688A1, 2016.
- [101] Takezawa, H.; Hayashi, M.; Iwasawa, Y.; Hosoi, M.; Iida, Y.; Tsuchiya, Y.; Horie, M.; Kamei, T. Substituted alkylamine derivatives. US5234946A, 1993.
- [102] Forster, M.; Chaikuad, A.; Bauer, Silke M.; Holstein, J.; Robers, Matthew B.; Corona, Cesear R.; Gehringer, M.; Pfaffenrot, E.; Ghoreschi, K.; Knapp, S.; Laufer, Stefan A. Selective JAK3 Inhibitors with a Covalent Reversible Binding Mode Targeting a New Induced Fit Binding Pocket. *Cell Chem. Biol.*, **2016**, *23* (11), 1335-1340.
- [103] Forster, M.; Chaikuad, A.; Dimitrov, T.; Döring, E.; Holstein, J.; Berger, B.-T.; Gehringer, M.; Ghoreschi, K.; Müller, S.; Knapp, S.; Laufer, S. A. Development, Optimization, and Structure-Activity Relationships of Covalent-Reversible JAK3 Inhibitors Based on a Tricyclic Imidazo[5,4-d]pyrrolo[2,3-b]pyridine Scaffold. *J. Med. Chem.*, **2018**, *61* (12), 5350-5366.
- [104] Hrast, M.; Rožman, K.; Jukič, M.; Patin, D.; Gobec, S.; Sova, M. Synthesis and structure-activity relationship study of novel quinazolinone-based inhibitors of MurA. *Bioorg. Med. Chem. Lett.*, **2017**, *27* (15), 3529-3533.
- [105] Snyder, D. S.; Tradtrantip, L.; Yao, C.; Kurth, M. J.; Verkman, A. S. Potent, Metabolically Stable Benzopyrimido-pyrrolo-oxazine-dione (BPO) CFTR Inhibitors for Polycystic Kidney Disease. *J. Med. Chem.*, **2011**, *54* (15), 5468-5477.

- [106] Sachdeva, N.; Dolzhenko, A. V.; Lim, S. J.; Ong, W. L.; Chui, W. K. An efficient synthesis of 2,4,7-trisubstituted pyrimido[1,2-a][1,3,5]triazin-6-ones. *New J. Chem.*, **2015**, 39 (6), 4796-4804.
- [107] Matasi, J. J.; Caldwell, J. P.; Hao, J.; Neustadt, B.; Arik, L.; Foster, C. J.; Lachowicz, J.; Tulshian, D. B. The discovery and synthesis of novel adenosine receptor (A2A) antagonists. *Bioorg. Med. Chem. Lett.*, **2005**, 15 (5), 1333-1336.
- [108] Bode, M. L.; Rousseau, A. L.; Gravestock, D.; Moleele, S. S.; Van der Westhuyzen, C. W. Imidazopyridines and imidazopyrimidines as HIV-1 reverse transcriptase inhibitors and their preparation, pharmaceutical compositions and use in the treatment of HIV infection. WO2010032195A1, 2010.
- [109] Fan, W.; Queneau, Y.; Popowycz, F. HMF in multicomponent reactions: utilization of 5-hydroxymethylfurfural (HMF) in the Biginelli reaction. *Green Chem.*, **2018**, 20 (2), 485-492.
- [110] Wu, Q.; Chen, J.; Guo, X.; Xu, Y. Copper(I)-Catalyzed Four-Component Coupling Using Renewable Building Blocks of CO<sub>2</sub> and Biomass-Based Aldehydes. *Eur. J. Org. Chem.*, **2018**, 2018 (24), 3105-3113.
- [111] Antonio, J. P. M.; Frade, R. F. M.; Santos, F. M. F.; Coelho, J. A. S.; Afonso, C. A. M.; Gois, P. M. P.; Trindade, A. F. NHC catalysed direct addition of HMF to diazo compounds: synthesis of acyl hydrazones with antitumor activity. *RSC Adv.*, **2014**, 4 (55), 29352-29356.
- [112] Baliani, A.; Bueno, G. J.; Stewart, M. L.; Yardley, V.; Brun, R.; Barrett, M. P.; Gilbert, I. H. Design and Synthesis of a Series of Melamine-based Nitroheterocycles with Activity against Trypanosomatid Parasites. *J. Med. Chem.*, **2005**, 48 (17), 5570-5579.
- [113] Kashiwagi, M.; Fuhshuku, K.-I.; Sugai, T. Control of the nitrile-hydrolyzing enzyme activity in *Rhodococcus rhodochrous* IFO 15564: preferential action of nitrile hydratase and amidase depending on the reaction condition factors and its application to the one-pot preparation of amides from aldehydes. *J. Mol. Catal. B: Enzym.*, **2004**, 29 (1), 249-258.
- [114] Ambreen, N.; Wirth, T. High-Temperature Synthesis of Amides from Alcohols or Aldehydes by Using Flow Chemistry. *Eur. J. Org. Chem.*, **2014**, 2014 (34), 7590-7593.
- [115] Jia, X.; Ma, J.; Wang, M.; Ma, H.; Chen, C.; Xu, J. Catalytic conversion of 5-hydroxymethylfurfural into 2,5-furandiamidine dihydrochloride. *Green Chem.*, **2016**, 18 (4), 974-978.
- [116] Cottier, L.; Descotes, G.; Soro, Y. Synthesis of Acetylated Ranunculin Diastereoisomers and  $\delta$ -Glucosyloxy- $\gamma$ -Oxo Esters from  $\alpha$  or  $\beta$  Glucosylmethylfurfural. *J. Carbohydr. Chem.*, **2005**, 24 (1), 55-71.
- [117] Casanova, O.; Iborra, S.; Corma, A. Biomass into chemicals: One pot-base free oxidative esterification of 5-hydroxymethyl-2-furfural into 2,5-dimethylfuroate with gold on nanoparticulated ceria. *J. Catal.*, **2009**, 265 (1), 109-116.
- [118] Kanai, S.; Nagahara, I.; Kita, Y.; Kamata, K.; Hara, M. A bifunctional cerium phosphate catalyst for chemoselective acetalization. *Chem. Sci.*, **2017**, 8 (4), 3146-3153.
- [119] Kim, M.; Su, Y.; Fukuoka, A.; Hensen, E. J. M.; Nakajima, K. Aerobic Oxidation of 5-(Hydroxymethyl)furfural Cyclic Acetal Enables Selective Furan-2,5-dicarboxylic Acid Formation with CeO<sub>2</sub>-Supported Gold Catalyst. *Angew. Chem. Int. Ed.*, **2018**, 57 (27), 8235-8239.
- [120] Arias, K. S.; Garcia-Ortiz, A.; Climent, M. J.; Corma, A.; Iborra, S. Mutual Valorization of 5-Hydroxymethylfurfural and Glycerol into Valuable Diol Monomers with Solid Acid Catalysts. *ACS Sustainable Chem. Eng.*, **2018**, 6 (3), 4239-4245.
- [121] Malleshham, B.; Sudarsanam, P.; Raju, G.; Reddy, B. M. Design of highly efficient Mo and W-promoted SnO<sub>2</sub> solid acids for heterogeneous catalysis: acetalization of bio-glycerol. *Green Chem.*, **2013**, 15 (2), 478-489.
- [122] Garcia-Ortiz, A.; Arias, K. S.; Climent, M. J.; Corma, A.; Iborra, S. One-Pot Synthesis of Biomass-Derived Surfactants by Reacting Hydroxymethylfurfural, Glycerol, and Fatty Alcohols on Solid Acid Catalysts. *ChemSusChem*, **2018**, 11 (17), 2870-2880.
- [123] Li, H.; Yang, T.; Riisager, A.; Saravanamurugan, S.; Yang, S. Chemoselective Synthesis of Dithioacetals from Bio-aldehydes with Zeolites under Ambient and Solvent-free Conditions. *ChemCatChem*, **2017**, 9 (6), 1097-1104.
- [124] Zhao, Q.; Zou, Y.; Huang, C.; Lan, P.; Zheng, J.; Ou, S. Formation of a Hydroxymethylfurfural-Cysteine Adduct and Its Absorption and Cytotoxicity in Caco-2 Cells. *J. Agric. Food Chem.*, **2017**, 65 (45), 9902-9908.
- [125] Mascal, M. 5-(Chloromethyl)furfural is the New HMF: Functionally Equivalent But More Practical in Terms of its Production From Biomass. *ChemSusChem*, **2015**, 8 (20), 3391-3395.
- [126] Sanda, K.; Rigal, L.; Gaset, A. Synthèse du 5-bromométhyl- et du 5-chlorométhyl-2-furanecarboxaldéhyde. *Carbohydr. Res.*, **1989**, 187 (1), 15-23.
- [127] Bredihhin, A.; Mäeorg, U.; Vares, L. Evaluation of carbohydrates and lignocellulosic biomass from different wood species as raw material for the synthesis of 5-bromomethylfurfural. *Carbohydr. Res.*,

- 2013**, 375, 63-67.
- [128] Romashov, L. V.; Ananikov, V. P. Synthesis of HIV-1 capsid protein assembly inhibitor (CAP-1) and its analogues based on a biomass approach. *Org. Biomol. Chem.*, **2016**, 14 (45), 10593-10598.
- [129] Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. The Vilsmeier Reaction: A New Synthetic Method for 5-(Chloromethyl)-2-furaldehyde. *Synthesis*, **1992**, 1992 (06), 541-542.
- [130] Newth, F. H.; Wiggins, L. F. 80. The conversion of sucrose into furan compounds. Part III. Some amidino-furans. *J. Chem. Soc.*, **1947**, 10.1039/JR9470000396 (0), 396-398.
- [131] Villain-Guillot, P.; Gualtieri, M.; Bastide, L.; Roquet, F.; Martinez, J.; Amblard, M.; Pugniere, M.; Leonetti, J.-P. Structure–Activity Relationships of Phenyl-Furanyl-Rhodanines as Inhibitors of RNA Polymerase with Antibacterial Activity on Biofilms. *J. Med. Chem.*, **2007**, 50 (17), 4195-4204.
- [132] Bedjeguelal, K.; Rabot, R.; Kaloun, E. B.; Mayer, P.; Marchand, A.; Rahier, N.; Schambel, P.; Bienayme, H. Preparation of pyrazolopyridine derivatives as ALK kinase inhibitors for treating cancer. WO2011045344A1, 2011.
- [133] Chundury, D.; Szmant, H. H. Preparation of polymeric building blocks from 5-hydroxymethyl- and 5-chloromethylfurfuraldehyde. *Ind. Eng. Chem. Prod. Res. Dev.*, **1981**, 20 (1), 158-163.
- [134] Larousse, C.; Rigal, L.; Gaset, A. Synthesis of 5,5'-oxydimethylenebis(2-furfural) by thermal dehydration of 5-hydroxymethyl-2-furfural in the presence of dimethylsulfoxide. *J. Chem. Technol. Biotechnol.*, **1992**, 53 (1), 111-116.
- [135] Cottier, L.; Descotes, G.; Eymard, L.; Rapp, K. Syntheses of  $\gamma$ -Oxo Acids or  $\gamma$ -Oxo Esters by Photooxygenation of Furanic Compounds and Reduction Under Ultrasound: Application to the Synthesis of 5-Aminolevulinic Acid Hydrochloride. *Synthesis*, **1995**, 1995 (03), 303-306.
- [136] Páez, A.; Rojas, H. A.; Portilla, O.; Sathicq, G.; Afonso, C. A. M.; Romanelli, G. P.; Martínez, J. J. Preyssler Heteropolyacids in the Self-Etherification of 5-Hydroxymethylfurfural to 5,5'-[Oxybis(methylene)]bis-2-furfural Under Mild Reaction Conditions. *ChemCatChem*, **2017**, 9 (17), 3322-3329.
- [137] Casanova, O.; Iborra, S.; Corma, A. Chemicals from biomass: Etherification of 5-hydroxymethyl-2-furfural (HMF) into 5,5'(oxy-bis(methylene))bis-2-furfural (OBMF) with solid catalysts. *J. Catal.*, **2010**, 275 (2), 236-242.
- [138] Wang, H.; Wang, Y.; Deng, T.; Chen, C.; Zhu, Y.; Hou, X. Carbocatalyst in biorefinery: Selective etherification of 5-hydroxymethylfurfural to 5,5'(oxy-bis(methylene))bis-2-furfural over graphene oxide. *Catal. Commun.*, **2015**, 59, 127-130.
- [139] Shinde, S.; Rode, C. Selective self-etherification of 5-(hydroxymethyl)furfural over Sn-Mont catalyst. *Catal. Commun.*, **2017**, 88, 77-80.
- [140] Mliki, K.; Trabelsi, M. Chemicals from biomass: Efficient and facile synthesis of 5,5'(oxy-bis(methylene))bis-2-furfural from 5-hydroxymethylfurfural. *Ind. Crops Prod.*, **2015**, 78, 91-94.
- [141] Wen, R.; Yu, F.; Dong, X.; Miao, Y.; Zhou, P.; Lin, Z.; Zheng, J.; Wang, H.; Huang, L.; Qing, D. Preparation of furfuryl compounds as antiviral agents. CN1456556A, 2003.
- [142] Wang, H.; Deng, T.; Wang, Y.; Cui, X.; Qi, Y.; Mu, X.; Hou, X.; Zhu, Y. Graphene oxide as a facile acid catalyst for the one-pot conversion of carbohydrates into 5-ethoxymethylfurfural. *Green Chem.*, **2013**, 15 (9), 2379.
- [143] Thombal, R. S.; Jadhav, V. H. Application of glucose derived magnetic solid acid for etherification of 5-HMF to 5-EMF, dehydration of sorbitol to isosorbide, and esterification of fatty acids. *Tetrahedron Lett.*, **2016**, 57 (39), 4398-4400.
- [144] Quiroz-Florentino, H.; Hernández-Benitez, R. I.; Aviña, J. A.; Burgueño-Tapia, E.; Tamariz, J. Total Synthesis of Naturally Occurring Furan Compounds 5-[[4-(4-Hydroxybenzyl)oxy]methyl]-2-furaldehyde and Pichiafuran C. *Synthesis*, **2011**, 2011 (07), 1106-1112.
- [145] Ilkei, V.; Faragó, K.; Sánta, Z.; Dékány, M.; Hazai, L.; Jr., C. S.; Szántay, C.; Kalas, G. The First Synthesis of Sessiline. *Int. J. Org. Chem.*, **2014**, 4, 309-313.
- [146] Berton, J. K. E. T.; Heugebaert, T. S. A.; Debrouwer, W.; Stevens, C. V. 3-Imidoallenylphosphonates: In Situ Formation and  $\beta$ -Alkoxylation. *Org. Lett.*, **2016**, 18 (2), 208-211.
- [147] El-Hajj, T.; Martin, J.-C.; Descotes, G. Dérivés de l'hydroxyméthyl-5 furfural. I. Synthèse de dérivés du di- et terfuranne. *J. Heterocycl. Chem.*, **1983**, 20 (1), 233-235.
- [148] de Freitas Filho, J. R.; Srivastava, R. M.; Soro, Y.; Cottier, L.; Descotes, G. Synthesis of new 2,3-unsaturated O-glycosides through Ferrier rearrangement. *J. Carbohydr. Chem.*, **2001**, 20 (7-8), 561-568.
- [149] Ding, Z.; Luo, X.; Ma, Y.; Chen, H.; Qiu, S.; Sun, G.; Zhang, W.; Yu, C.; Wu, Z.; Zhang, J. Eco-friendly synthesis of 5-hydroxymethylfurfural (HMF) and its application to the Ferrier-rearrangement reaction. *J. Carbohydr. Chem.*, **2018**, 37 (2), 81-93.
- [150] Jogia, M.; Vakamoce, V.; Weavers, R. Synthesis of Some Furfural and Syringic Acid Derivatives. *Aust. J. Chem.*, **1985**, 38 (7), 1009-1016.
- [151] Bognár, R.; Herczegh, P.; Zsély, M.; Batta, G. Synthesis of 3,4-dideoxy-dl-hex-3-enopyranosides from

- 5-hydroxymethyl-2-furaldehyde. *Carbohydr. Res.*, **1987**, *164*, 465-469.
- [152] Bakholdina, L. A.; Khlebnikov, A. I.; Sevodin, V. P. Mild reaction of primary alcohols with ferulic acid. *Russ. J. Org. Chem.*, **2016**, *52* (3), 441-443.
- [153] Zhou, H.; Liu, W.; Sun, C.; Peng, C.; Wang, J.; Wang, Q.; Yang, C. Synthesis of Novel Coumarin Derivatives and in vitro Biological Evaluation as Potential PTP 1B Inhibitors. *Heterocycles*, **2013**, *87* (8), 1711.
- [154] Quiroz-Florentino, H.; García, A.; Burgueño-Tapia, E.; Tamariz, J. Total synthesis of the natural succinate derivative of 5-(hydroxymethyl)furfural isolated from the Noni fruit (*Morinda citrifolia*). *Nat. Prod. Res.*, **2009**, *23* (14), 1355-1362.
- [155] Sugimura, H.; Kikuchi, M.; Kato, S.; Sekita, W.; Sasaki, I. Practical synthesis of mume-fural, a component of Japanese apricot juice concentrate. *Tetrahedron*, **2016**, *72* (47), 7638-7641.
- [156] Krystof, M.; Pérez-Sánchez, M.; Domínguez de María, P. Lipase-Catalyzed (Trans)esterification of 5-Hydroxy- methylfurfural and Separation from HMF Esters using Deep-Eutectic Solvents. *ChemSusChem*, **2013**, *6* (4), 630-634.
- [157] Qin, Y.-Z.; Zong, M.-H.; Lou, W.-Y.; Li, N. Biocatalytic Upgrading of 5-Hydroxymethylfurfural (HMF) with Levulinic Acid to HMF Levulinate in Biomass-Derived Solvents. *ACS Sustainable Chem. Eng.*, **2016**, *4* (7), 4050-4054.
- [158] Dow, R. L.; Kelly, R. C.; Schletter, I.; Wierenga, W. A Direct Alcohol for Hydrazine Interchange: Scope and Stereochemistry. *Synth. Commun.*, **1981**, *11* (1), 43-53.
- [159] Lewis, T. A.; Bayless, L.; Eckman, J. B.; Ellis, J. L.; Grewal, G.; Libertine, L.; Marie Nicolas, J.; Scannell, R. T.; Wels, B. F.; Wenberg, K.; Wypij, D. M. 5-Lipoxygenase inhibitors with histamine H1 receptor antagonist activity. *Bioorg. Med. Chem. Lett.*, **2004**, *14* (9), 2265-2268.
- [160] Iovel, I.; Mertins, K.; Kischel, J.; Zapf, A.; Beller, M. An Efficient and General Iron-Catalyzed Arylation of Benzyl Alcohols and Benzyl Carboxylates. *Angew. Chem. Int. Ed.*, **2005**, *44* (25), 3913-3917.
- [161] Zhou, X.; Rauchfuss, T. B. Production of Hybrid Diesel Fuel Precursors from Carbohydrates and Petrochemicals Using Formic Acid as a Reactive Solvent. *ChemSusChem*, **2013**, *6* (2), 383-388.
- [162] Bering, L.; Jeyakumar, K.; Antonchick, A. P. Metal-Free C–O Bond Functionalization: Catalytic Intramolecular and Intermolecular Benzoylation of Arenes. *Org. Lett.*, **2018**, *20* (13), 3911-3914.
- [163] Onorato, A.; Pavlik, C.; Invernale, M. A.; Berghorn, I. D.; Sotzing, G. A.; Morton, M. D.; Smith, M. B. Polymer-mediated cyclodehydration of alditols and ketohexoses. *Carbohydr. Res.*, **2011**, *346* (13), 1662-1670.
- [164] Nale, S. D.; Jadhav, V. H. Synthesis of Fuel Intermediates from HMF/Fructose. *Catal. Lett.*, **2016**, *146* (10), 1984-1990.
- [165] Ryabukhin, D. S.; Zakusilo, D. N.; Kompanets, M. O.; A.Tarakanov, A.; Boyarskaya, I. A.; Artamonova, T. O.; Khohodorkovskiy, M. A.; Opeida, I. O.; Vasilyev, A. V. Superelectrophilic activation of 5-hydroxymethylfurfural and 2,5-diformylfuran: organic synthesis based on biomass-derived products. *Beilstein J. Org. Chem.*, **2016**, *12*, 2125-2135.
- [166] Teunissen, H. P. Velocity measurements on the opening of the furane ring in hydroxymethylfurfuraldehyde. *Recl. Trav. Chim. Pays-Bas*, **1930**, *49* (9), 784-826.
- [167] Luijckx, G. C. A.; van Rantwijk, F.; van Bekkum, H. Hydrothermal formation of 1,2,4-benzenetriol from 5-hydroxymethyl-2-furaldehyde and d-fructose. *Carbohydr. Res.*, **1993**, *242*, 131-139.
- [168] Tšupova, S.; Rominger, F.; Rudolph, M.; Hashmi, A. S. K. Synthesis of phenols from hydroxymethylfurfural (HMF). *Green Chem.*, **2016**, *18* (21), 5800-5805.
- [169] Roylance, J. J.; Choi, K.-S. Electrochemical reductive biomass conversion: direct conversion of 5-hydroxymethylfurfural (HMF) to 2,5-hexanedione (HD) via reductive ring-opening. *Green Chem.*, **2016**, *18* (10), 2956-2960.
- [170] Ren, D.; Song, Z.; Li, L.; Liu, Y.; Jin, F.; Huo, Z. Production of 2,5-hexanedione and 3-methyl-2-cyclopenten-1-one from 5-hydroxymethylfurfural. *Green Chem.*, **2016**, *18* (10), 3075-3081.
- [171] Wu, W.-P.; Xu, Y.-J.; Zhu, R.; Cui, M.-S.; Li, X.-L.; Deng, J.; Fu, Y. Selective Conversion of 5-Hydroxymethylfuraldehyde Using Cp\*Ir Catalysts in Aqueous Formate Buffer Solution. *ChemSusChem*, **2016**, *9* (10), 1209-1215.
- [172] Xu, Z.; Yan, P.; Li, H.; Liu, K.; Liu, X.; Jia, S.; Zhang, Z. C. Active Cp\*Iridium(III) Complex with ortho-Hydroxyl Group Functionalized Bipyridine Ligand Containing an Electron-Donating Group for the Production of Diketone from 5-HMF. *ACS Catal.*, **2016**, *6* (6), 3784-3788.
- [173] Wozniak, B.; Li, Y. H.; Hinze, S.; Tin, S.; de Vries, J. G. Efficient Synthesis of Biomass-Derived N-Substituted 2-Hydroxymethyl-5-Methyl-Pyrroles in Two Steps from 5-Hydroxymethylfurfural. *Eur. J. Org. Chem.*, **2018**, *2018* (17), 2009-2012.
- [174] Sacia, E. R.; Deaner, M. H.; Louie, Y. L.; Bell, A. T. Synthesis of biomass-derived methylcyclopentane as a gasoline additive via aldol condensation/hydrodeoxygenation of 2,5-hexanedione. *Green Chem.*,

- 2015**, *17* (4), 2393-2397.
- [175] Duan, Y.; Zheng, M.; Li, D.; Deng, D.; Ma, L.-F.; Yang, Y. Conversion of HMF to methyl cyclopentenolone using Pd/Nb<sub>2</sub>O<sub>5</sub> and Ca–Al catalysts via a two-step procedure. *Green Chem.*, **2017**, *19* (21), 5103-5113.
- [176] Wozniak, B.; Spannenberg, A.; Li, Y.; Hinze, S.; de Vries, J. G. Cyclopentanone Derivatives from 5-Hydroxymethylfurfural via 1-Hydroxyhexane-2,5-dione as Intermediate. *ChemSusChem*, **2018**, *11* (2), 356-359.
- [177] Ohyama, J.; Kanao, R.; Ohira, Y.; Satsuma, A. The effect of heterogeneous acid-base catalysis on conversion of 5-hydroxymethylfurfural into a cyclopentanone derivative. *Green Chem.*, **2016**, *18* (3), 676-680.
- [178] Ohyama, J.; Kanao, R.; Esaki, A.; Satsuma, A. Conversion of 5-hydroxymethylfurfural to a cyclopentanone derivative by ring rearrangement over supported Au nanoparticles. *Chem. Commun.*, **2014**, *50* (42), 5633-5636.
- [179] Verrier, C.; Moebs-Sanchez, S.; Queneau, Y.; Popowycz, F. The Piancatelli reaction and its variants: recent applications to high added-value chemicals and biomass valorization. *Org. Biomol. Chem.*, **2018**, *16* (5), 676-687.
- [180] Lichtenhaler, F. W.; Brust, A.; Cuny, E. Sugar-derived building blocks. Part 26. Hydrophilic pyrroles, pyridazines and diazepinones from -fructose and isomaltulose. *Green Chem.*, **2001**, *3* (5), 201-209.
- [181] Heugebaert, T. S. A.; Stevens, C. V.; Kappe, C. O. Singlet-Oxygen Oxidation of 5-Hydroxymethylfurfural in Continuous Flow. *ChemSusChem*, **2015**, *8* (10), 1648-1651.
- [182] Kucherov, F. A.; Galkin, K. I.; Gordeev, E. G.; Ananikov, V. P. Efficient route for the construction of polycyclic systems from bioderived HMF. *Green Chem.*, **2017**, *19* (20), 4858-4864.
- [183] Galkin, K. I.; Kucherov, F. A.; Markov, O. N.; Egorova, K. S.; Posvyatenko, A. V.; Ananikov, V. P. Facile Chemical Access to Biologically Active Norcantharidin Derivatives from Biomass. *Molecules*, **2017**, *22* (12), 2210.
- [184] Higson, S.; Subrizi, F.; Sheppard, T. D.; Hailes, H. C. Chemical cascades in water for the synthesis of functionalized aromatics from furfurals. *Green Chem.*, **2016**, *18* (7), 1855-1858.
- [185] Lin, Z.; Ierapetritou, M.; Nikolakis, V. Aromatics from Lignocellulosic Biomass: Economic Analysis of the Production of p-Xylene from 5-Hydroxymethylfurfural. *AIChE J.*, **2013**, *59* (6), 2079-2087.
- [186] Ni, L.; Xin, J.; Dong, H.; Lu, X.; Liu, X.; Zhang, S. A Simple and Mild Approach for the Synthesis of p-Xylene from Bio-Based 2,5-Dimethylfuran by Using Metal Triflates. *ChemSusChem*, **2017**, *10* (11), 2394-2401.
- [187] Shiramizu, M.; Toste, F. D. On the Diels–Alder Approach to Solely Biomass-Derived Polyethylene Terephthalate (PET): Conversion of 2,5-Dimethylfuran and Acrolein into p-Xylene. *Chem. Eur. J.*, **2011**, *17* (44), 12452-12457.
- [188] Serum, E. M.; Sermadurai, S.; Zimmermann, N.; Sibi, M. P. Valorization of 2,5-Furandicarboxylic Acid. Diels-Alder Reactions with Benzyne. *Green Chem.*, **2018**, *20* (7), 1448-1454.
- [189] Pacheco, J. J.; Labinger, J. A.; Sessions, A. L.; Davis, M. E. Route to Renewable PET: Reaction Pathways and Energetics of Diels–Alder and Dehydrative Aromatization Reactions Between Ethylene and Biomass-Derived Furans Catalyzed by Lewis Acid Molecular Sieves. *ACS Catal.*, **2015**, *5* (10), 5904-5913.
- [190] Chang, C.-C.; Je Cho, H.; Yu, J.; Gorte, R. J.; Gulbinski, J.; Dauenhauer, P.; Fan, W. Lewis acid zeolites for tandem Diels-Alder cycloaddition and dehydration of biomass-derived dimethylfuran and ethylene to renewable p-xylene. *Green Chem.*, **2016**, *18* (5), 1368-1376.
- [191] Song, S.; Wu, G.; Dai, W.; Guan, N.; Li, L. Diels-Alder and dehydration reactions of furan derivatives with ethylene catalyzed by liquid Brønsted acids and Lewis acids. *J. Mol. Catal. A: Chem.*, **2016**, *420*, 134-141.
- [192] Do, P. T. M.; McAtee, J. R.; Watson, D. A.; Lobo, R. F. Elucidation of Diels–Alder Reaction Network of 2,5-Dimethylfuran and Ethylene on HY Zeolite Catalyst. *ACS Catal.*, **2013**, *3* (1), 41-46.
- [193] Pacheco, J. J.; Davis, M. E. Synthesis of terephthalic acid via Diels-Alder reactions with ethylene and oxidized variants of 5-hydroxymethylfurfural. *Proc. Natl. Acad. Sci. U. S. A.*, **2014**, *111* (23), 8363-8367.
- [194] McNelis, B. J.; Sternbach, D. D.; MacPhail, A. T. Synthetic and kinetic studies of substituent effects in the furan intramolecular Diels-Alder reaction. *Tetrahedron*, **1994**, *50* (23), 6767-6782.
- [195] Sun, S.; Murray, W. V. Solid Phase Diels–Alder Reactions of Amino Acid Derived Trienes. *J. Org. Chem.*, **1999**, *64* (16), 5941-5945.
- [196] Schinzer, D.; Bourguet, E.; Ducki, S. Synthesis of Furano-Epothilone D. *Chem. Eur. J.*, **2004**, *10* (13), 3217-3224.
- [197] Koh, P. F.; Loh, T. P. Synthesis of biologically active natural products, aspergillides A and B, entirely from biomass derived platform chemicals. *Green Chem.*, **2015**, *17* (7), 3746-3750.
- [198] Connolly, T. J.; Considine, J. L.; Ding, Z.; Forsatz, B.; Jennings, M. N.; MacEwan, M. F.; McCoy, K.

- M.; Place, D. W.; Sharma, A.; Sutherland, K. Efficient Synthesis of 8-Oxa-3-aza-bicyclo[3.2.1]octane Hydrochloride. *Org. Process Res. Dev.*, **2010**, *14* (2), 459-465.
- [199] Oikawa, M.; Ikoma, M.; Sasaki, M. Parallel synthesis of tandem Ugi/Diels–Alder reaction products on a soluble polymer support directed toward split-pool realization of a small molecule library. *Tetrahedron Lett.*, **2005**, *46* (3), 415-418.
- [200] Gupta, P.; Singh, S. K.; Pathak, A.; Kundu, B. Template-directed approach to solid-phase combinatorial synthesis of furan-based libraries. *Tetrahedron*, **2002**, *58* (52), 10469-10474.
- [201] Müller, C.; Diehl, V.; Lichtenthaler, F. W. Building blocks from sugars. Part 23. Hydrophilic 3-pyridinols from fructose and isomaltulose. *Tetrahedron*, **1998**, *54* (36), 10703-10712.
- [202] Villard, R.; Robert, F.; Blank, I.; Bernardinelli, G.; Soldo, T.; Hofmann, T. Racemic and Enantiopure Synthesis and Physicochemical Characterization of the Novel Taste Enhancer N-(1-Carboxyethyl)-6-(hydroxymethyl)pyridinium-3-ol Inner Salt. *J. Agric. Food Chem.*, **2003**, *51* (14), 4040-4045.
- [203] Koch, J.; Pischetsrieder, M.; Polborn, K.; Severin, T. Formation of pyridinium betaines by reaction of hexoses with primary amines. *Carbohydr. Res.*, **1998**, *313* (2), 117-123.
- [204] Frank, O.; Ottinger, H.; Hofmann, T. Characterization of an Intense Bitter-Tasting 1H,4H-Quinolizinium-7-olate by Application of the Taste Dilution Analysis, a Novel Bioassay for the Screening and Identification of Taste-Active Compounds in Foods. *J. Agric. Food Chem.*, **2001**, *49* (1), 231-238.
- [205] Soldo, T.; Hofmann, T. Application of Hydrophilic Interaction Liquid Chromatography/Comparative Taste Dilution Analysis for Identification of a Bitter Inhibitor by a Combinatorial Approach Based on Maillard Reaction Chemistry. *J. Agric. Food Chem.*, **2005**, *53* (23), 9165-9171.
- [206] Sowmiah, S.; Veiros, L. F.; Esperanca, J. M.; Rebelo, L. P.; Afonso, C. A. Organocatalyzed One-Step Synthesis of Functionalized N-Alkyl-Pyridinium Salts from Biomass Derived 5-Hydroxymethylfurfural. *Org. Lett.*, **2015**, *17* (21), 5244-5247.
- [207] Rajmohan, R.; Gayathri, S.; Vairaprakash, P. Facile synthesis of 5-hydroxymethylfurfural: a sustainable raw material for the synthesis of key intermediates toward 21,23-dioxaporphyrins. *RSC Adv.*, **2015**, *5* (121), 100401-100407.
- [208] McDermott, P. J.; Stockman, R. A. Combining Two-Directional Synthesis and Tandem Reactions: Synthesis of Trioxadispiroketal. *Org. Lett.*, **2005**, *7* (1), 27-29.